## Recommended Routine Immunization Schedule for Infants, Children, and Adults



|   | Age  |             |             |              |              |              |            |                 |                |               |
|---|--|-------------|-------------|--------------|--------------|--------------|------------|-----------------|----------------|---------------|
| Vaccine   | 2<br>months  | 4<br>months | 6<br>months | 12<br>months | 18<br>months | 4-6<br>years | Grade<br>6 | Grade<br>8 or 9 | All<br>adults  | 65<br>years   |
| Diphtheria, Tetanus, Pertussis,<br>Polio, Haemophilus influenzae<br>type b (DTaP-IPV-Hib) | •  | *           | *           |              | •            |              |            |                 |                |               |
| Pneumococcal conjugate<br>15-valent (Pneu-C-15)*  | •  | •           |             | •            |              |              |            |                 |                |               |
| Rotavirus   | *  | •           |             |              |              |              |            |                 |                |               |
| Measles, Mumps, Rubella,<br>Varicella (MMRV)  |  |             |             | •            |              | •            |            |                 |                |               |
| Tetanus, Diphtheria, Pertussis,<br>Polio (Tdap-IPV)                                       |  |             |             |              |              | •            |            |                 |                |               |
| Hepatitis B (HB)  |  |             |             |              |              |              | **         |                 |                |               |
| Human Papillomavirus (HPV)  |  |             |             |              |              |              | **         |                 |                |               |
| Meningococcal<br>Conjugate Quadrivalent<br>(Men-C-ACYW)                                   |  |             |             | <b>*</b>     |              |              | <b>*</b>   |                 |                |               |
| Tetanus, Diphtheria,<br>Pertussis (Tdap)  |  |             |             |              |              |              |            | •               | •              |               |
| Tetanus, Diphtheria (Td)  |  |             |             |              |              |              |            |                 | Every 10 years |               |
| Pneumococcal conjugate<br>20-valent (Pneu-C-20)   |  |             |             |              |              |              |            |                 |                | Lifetime dose |
| Influenza (Flu)   | All Manitobans 6 months of age and older are eligible for influenza vaccine each year. People aged 65 years and older are eligible for the high-dose influenza vaccine |             |             |              |              |              |            |                 |                |               |

- ♦ = A single vaccine dose given.
- Tdap vaccine is available for adults who are due for their tetanus booster and have not had a pertussis-containing vaccine in adulthood and pregnant women in every pregnancy between 27 and 32 weeks gestation.
- \* Children with high risk medical conditions and those living in First Nations communities are eligible to receive four doses of the PCV20 vaccine at 2, 4, 6 and 18 months.

A different schedule may be needed for high risk individuals, or if one or more doses of a vaccine are missed. For more information about immunization, talk to your health-care provider or visit:

www.manitoba.ca/health/publichealth/cdc/div/index.html