

Vaccine Clinic Resource for Immunizers

Disclaimer: *this Quick Reference is not intended to replace other product specific vaccine references. The document is intended as a quick reference for frequently referred to information. Please refer to the product monograph and vaccine specific resources for all current and complete information.*

Title:	School Based Clinic (Grade 6 and Grade 8/9) - Quick Reference Guide <i>This guide also provides additional information for those who may have missed the school based clinics and require catch-up immunizations.</i>
Effective Date:	July 27, 2023
Approver:	Final

School Based Vaccine Resources:

Fact sheets and Product Monographs

<https://www.manitoba.ca/health/publichealth/cdc/div/sip.html>

Eligibility Criteria

For the most up to date information on eligibility criteria refer to <https://www.manitoba.ca/health/publichealth/cdc/vaccineeligibility.html>

Manitoba Immunization Schedule:

<https://www.manitoba.ca/health/publichealth/cdc/div/schedules.html>

Canadian Immunization Guide:

For additional guidance on contraindications, precautions and special populations refer to each vaccine specific section:

<https://www.canada.ca/en/public-health/services/publications/healthy-living/canadian-immunization-guide-part-4-active-vaccines.html>

Hepatitis B Vaccine (HBV)

Product	Storage and Handling	Eligibility Criteria	Recommendations for Use
<p>ENGERIX®-B Hepatitis B vaccine (recombinant)</p> <p>Format: Pre-filled 1.0 mL syringe (single dose) or Pre-filled 0.5 mL syringe (single dose) (1 per/box)</p> <p><i>RECOMBIVAX HB® had been used routinely in Manitoba in the past and can be interchanged with ENGERIX®-B NACI Recommendation - Monovalent hepatitis B vaccines may be used interchangeably, according to the recommended dosage and schedule.</i></p>	<p>Storage: Refrigerated 2° to 8°C until expiry date on label.</p> <p>Do NOT freeze. Protect from light.</p> <p>Handling: Reconstitution NOT required</p> <p>The vaccine should be inspected visually for any foreign particulate matter and/or coloration prior to administration.</p> <p>Before use of ENGERIX-B, the vaccine should be well shaken to resuspend the sediment of fine white particles of adjuvant (aluminum hydroxide) which settles during storage and to obtain a slightly opaque, white suspension. Discard if the content appears otherwise.</p>	<ul style="list-style-type: none"> All children born on or after January 1st, 2006 in the grade 6* school-based program that are 11-15** years of age are eligible for the 1.0ml, 2 dose schedule <p>*Children who are 10 years of age for their first dose of HBV in Grade 6 (birthday between Sept and Dec of that year), are eligible for the 1.0 ml, 2 dose schedule.</p> <p>**Children 11 to 15 years of age who have had one or two pediatric doses (0.5mL/dose) will require 3 doses to complete their series. They can complete with the 0.5 ml (10mcg) dose(s), or if necessary, immunizers can finish the 3-dose series with 1.0 mL (20mcg) dose(s) (e.g. vaccine supply available is prefilled 1.0 mL syringe only).</p> <ul style="list-style-type: none"> Children/adolescents 16-18 years of age Individuals of any age who meet the high-risk criteria may be eligible for 3 or 4 doses of hepatitis B vaccine. Refer to Manitoba Health Eligibility Criteria and Canadian Immunization Guide for eligibility and dosage requirements 	<p>Regimen: 2 dose schedule: 0, 6 months</p> <p>Dosage: 1.0 ml (20 mcg)</p> <p>Route: IM</p> <p>Interval: Minimum interval is 6 months (24 weeks)</p> <p><i>Note re: PHIMs forecaster:</i></p> <ul style="list-style-type: none"> If both 1.0ml doses are not received before the age of 16, the forecast will revert to a three-dose schedule. If the 2nd dose is less than the minimum interval it will forecast for 3 doses If the dosage of the Hep B product is not specified or if the dosage is < 1 ml it will forecast a 3 dose 0.5 ml/dose schedule <p>Regimen: 3 dose schedule: 0, 1, 6 months</p> <p>Dosage: 0.5 ml (10 mcg)</p> <p>Route: IM</p> <p>Interval: Although a schedule of months 0, 1 and at least 2 is authorized, the preferred schedule is months 0, 1 and 6</p>

Product is latex and preservative/thimerosal free

Potential Allergens: Yeast protein **Other Ingredients:** aluminum hydroxide (adjuvant), disodium phosphate dehydrate, sodium dihydrogen phosphate dehydrate, sodium chloride, water for injection

Pregnancy and lactation: Can be used safely during pregnancy and breastfeeding

Human Papillomavirus – 9 (HPV-9) vaccine			
Product	Storage and Handling	Eligibility Criteria	Recommendations for Use
<p>Gardasil®9 Human Papillomavirus 9-valent Vaccine, Recombinant</p> <p>HPV (6, 11,16,18,31,33, 45,52, and 58) L1 protein</p> <p>Format: Pre-filled 0.5 mL syringe (10 per/box)</p> <p>Whenever possible, the same HPV vaccine should be used to complete a vaccine series. However, if previously immunized with dose(s) of Gardasil®4 then complete the series with Gardasil®9 (CIG).</p>	<p>Storage: Refrigerated 2° to 8°C until expiry date on label.</p> <p>Do NOT freeze. Protect from light.</p> <p>Handling: Reconstitution NOT required</p> <p>Gardasil®9 should be administered as soon as possible after being removed from refrigeration.</p> <p>Shake well before use. Thorough agitation immediately before administration is necessary to maintain suspension of the vaccine.</p> <p>Discard the product if it is frozen, particulates are present, or if it appears discolored.</p>	<ul style="list-style-type: none"> Individuals (immunocompetent) 9 to 14 years of age are eligible to receive the HPV vaccine, routinely provided as part of the grade 6 school-based immunization program. 	<p>Regimen: 2 dose schedule: 0, 6 months</p> <p>Dosage: 0.5 mL</p> <p>Route: IM</p> <p>Interval: Minimum interval between the first and second dose of vaccine is 6 months (24 weeks). If a second dose is administered earlier than 6 months, a third dose should be given at least 6 months (24 weeks) after the first dose.</p>
		<ul style="list-style-type: none"> Individuals (immunocompetent) 15 years* of age and older if born on or after 1997 (females) OR born on or after 2002 (males) <i>*For those who missed the school-based program, a 3-dose schedule is recommended, unless the first dose of vaccine was administered before the age of 15 yrs.</i> 	<p>Regimen: 3 dose schedule: 0, 2, 6 months</p> <p>Dosage: 0.5 mL</p> <p>Route: IM</p> <p>Interval: The minimum interval between the first and second doses of vaccine is 4 weeks (1 month), the minimum interval between the second and third doses of vaccine is 12 weeks (3 months), and the minimum interval between the first and last doses is 24 weeks (6 months).</p>
		<ul style="list-style-type: none"> Individuals 9 years and older who meet the high-risk criteria may be eligible for 3 doses of HPV vaccine. Refer to Manitoba Health Eligibility Criteria and Canadian Immunization Guide for eligibility and dosage requirements 	
<p>Product is latex and preservative/thimerosal free</p> <p>Potential Allergens: Yeast protein, polysorbate 80 (may also be found in bowel preparation, PEG, laxatives, corn syrup, cosmetics, etc.) Other Ingredients: aluminum (as amorphous aluminum hydroxyphosphate sulfate adjuvant), L-histidine, sodium borate, sodium chloride, water for injection</p> <p>Pregnancy and lactation: HPV vaccines are not recommended for use in pregnancy because data on HPV vaccination in pregnancy are limited. If a woman is found to be pregnant after initiating the vaccination series, completion of the series should be delayed until after pregnancy. No intervention is required if vaccine has been administered during pregnancy. Limited data on breastfeeding, but not contraindicated.</p>			

Meningococcal Vaccine (Quadrivalent)			
Product	Storage and Handling	Eligibility Criteria	Recommendations for Use
<p>Menactra®</p> <p>Meningococcal (Groups A,C,Y and W-135) Polysaccharide Diphtheria Toxoid Conjugate Vaccine</p> <p>Format: Single and 5/per box 0.5ml vials</p>	<p>Storage: Refrigerated 2°to 8°C until expiry date on label.</p> <p>Do NOT freeze.</p> <p>Handling: Reconstitution NOT required Shake the vaccine well until a uniform, clear to slightly turbid liquid results.</p>	<ul style="list-style-type: none"> Individuals born during or after January 1, 2008 are eligible to receive one dose during the Grade 6 school based program. 	<p>Regimen: 1 dose*</p> <p>Dosage: 0.5 mL</p> <p>Route: IM</p> <p>Interval: <i>*Those with certain high-risk medical conditions may require additional doses see Manitoba Health eligibility criteria for complete list and refer to Canadian Immunization guide for dose requirements and intervals</i></p>
<p>Product is latex and preservative/thimerosal free.</p> <p>Potential Allergens: Diphtheria toxoid carrier protein Other Ingredients: Sodium chloride, sodium phosphate dibasic anhydrous, sodium phosphate monobasic; water for injection</p> <p>Pregnancy and lactation: Conjugate quadrivalent meningococcal vaccine should be considered in pregnancy. Limited data on breastfeeding, but not contraindicated.</p>			
Product	Storage and Handling	Eligibility Criteria	Recommendations for Use
<p>Nimenrix®</p> <p>Meningococcal polysaccharide groups A, C, W-135 and Y conjugate vaccine</p> <p>Format: Single dose vial of sterile lyophilized white powder or cake; diluent presented in a pre-filled syringe (0.5 ml)</p> <p>Single or 10/box vial</p>	<p>Storage: Refrigerated 2°to 8°C until expiry date on label.</p> <p>Do NOT freeze. Protect from light.</p> <p>Handling: <u>Reconstitution required</u> Add the entire contents of the pre-filled syringe of diluent to the vial containing the powder. The reconstituted vaccine is a clear, colourless solution. After reconstitution, the vaccine should be used promptly.</p>	<ul style="list-style-type: none"> Individuals born during or after January 1, 2008 are eligible to receive one dose during the Grade 6 school based program. 	<p>Regimen: 1 dose*</p> <p>Dosage: 0.5 mL</p> <p>Route: IM</p> <p>Interval: <i>*Those with certain high-risk medical conditions may require additional doses - see Manitoba Health eligibility criteria for complete list and refer to the Canadian Immunization guide for dose requirements and intervals</i></p>
<p>Product is latex and preservative/thimerosal free.</p> <p>Potential Allergens: Tetanus toxoid carrier protein Other Ingredients: Sucrose, Trometamol, Sodium chloride, water for injection</p> <p>Pregnancy and lactation: Conjugate quadrivalent meningococcal vaccine should be considered in pregnancy. Limited data on breastfeeding, but not contraindicated.</p>			

Tetanus, Diphtheria, acellular Pertussis (Tdap) Vaccine			
Product	Storage and Handling	Eligibility Criteria	Recommendations for Use
Boostrix® Combined diphtheria, tetanus, acellular pertussis (adsorbed) vaccine for booster vaccination Format: Pre-filled 0.5 mL syringe 10 per/box	Storage: Refrigerated 2° to 8°C until expiry date on label. Do not freeze. Protect from light. Handling: Reconstitution NOT required Shake vaccine well to obtain a homogeneous turbid white liquid.	<ul style="list-style-type: none"> Individuals 13 to 15 years of age (Grade 8 or 9 school-based program). <p><i>Any dose after the age of 10 can be counted as an adolescent booster.</i></p>	Regimen: 1 dose* Dosage: 0.5 mL Route: IM Interval: There is no minimum interval between the Td and Tdap vaccine when Tdap is being given for pertussis protection (NACI).
Product is latex and preservative/thimerosal free Potential Allergens: polysorbate 80 (may also be found in bowel preparation, PEG, laxatives, corn syrup, cosmetics, etc.) Other Ingredients: Formaldehyde, glycerin, sodium chloride, water for injection, aluminum (as aluminum salts) Pregnancy and lactation: Can be used safely during pregnancy and breastfeeding			

Tetanus, Diphtheria, acellular Pertussis and Polio (Tdap-IPV) Vaccine			
Product	Storage and Handling	Eligibility Criteria	Recommendations for Use
Boostrix-Polio® Combined diphtheria, tetanus, acellular pertussis (adsorbed) and inactivated poliomyelitis vaccine for booster vaccination Format: Pre-filled 0.5 mL syringe 10 per/box	Storage: Refrigerated 2° to 8°C until expiry date on label. Do not freeze. Protect from light. Handling: Reconstitution NOT required Shake vaccine well to obtain a homogeneous turbid white liquid.	<ul style="list-style-type: none"> Individuals 7-17 years of age that also require the polio antigen (IPV) 	Regimen: 1 dose* *Some students may be forecasted for additional catch-up doses. Dosage: 0.5 mL Route: IM Interval: Refer to the following table for dose requirements and intervals: https://www.manitoba.ca/health/publichealth/cdc/div/not.html#more
Product is latex and preservative/thimerosal free Potential Allergens: neomycin, polymycin B sulphate Other Ingredients: Aluminum (as aluminum salts), formaldehyde, medium 199, sodium chloride, water for injection Pregnancy and lactation: Can be used during pregnancy and breastfeeding			