

# Progress of geological mapping at Paint Lake, Manitoba: identification of a new metasedimentary succession in the Thompson Nickel Belt



C.G. Couëslan

#### Summary

The Paint Lake area is underlain by predominantly multicomponent gneiss with significant exposures of enderbitic gneiss and layered mafic rocks along the west side of the lake. A sequence of possibly Archean supracrustal rocks, consisting dominantly of metagreywacke and metapsammite with minor iron formation, metapelite, and metavolcanic rocks, is present as disrupted bands along the east and west shores, and a relatively continuous belt through the central islands of the lake. A variably metasomatized peralkaline melasyenite intrusion with dykes of alkali feldspar granite has been identified in the south west corner of the lake. A carbonatite dyke swarm identified in the previous year has been extended along strike for over 21 km and may continue north of the currently mapped area. The dominant structures in the area consist of upright isoclinal, shallowly to moderately plunging folds that have deformed the regional foliation. Significant shear zones are manifested by metre-scale zones of mylonite to ultramylonite along the east shore of the lake.

Similarities of the supracrustal rocks at Paint Lake with rocks of the Ospwagan Group

1) extensive psammitic horizons which could be misinterpreted as Manasan Formation M1 quartzite or Setting Formation quartzite;


2) siliceous, calcareous units which could be confused with certain horizons of Thompson Formation;

3) sillimanite- and garnet-rich horizons of metapelite similar to Pipe Formation P2 member; 4) silicate facies iron formation similar to Pipe Formation;


5) concretion-bearing quartzite with interbedded metapelite similar to Setting Formation. The Archean supracrustal rocks are likely derived from a more juvenile source than similar

rocks of the Ospwagan Group. They are generally depleted in incompatible large ion lithophile elements such as Cs, Rb, K, and U, and enriched in Sr. This is likely a function of a difference in provenance between the two metasedimentary groups. The Archean metasediments are most likely the product of detritus derived from an arc/greenstone belt. The Ospwagan Group is most likely sourced from the peneplanation of the Superior craton which consists largely of Archean granitoid rocks.


## 1) Comparison of Archean supracrustal and Ospwagan Group lithologies























k) Archean interbedded metapsammite and m) Garnet-rich calcareous concretions in Archean metapsammite. A thin pelitic layer is visible at the bottom of the photo. n) Garnet-rich calcareous concretion in Setting Formation quartzite at Mystery Lake. A thin pelitic layer is visible at the top of the photo. Cnc = concretion.

# Paint Lake Geology



metagreywacke.b) Manasan Formation, M1 quartzite in contact with Manasan Formation M2 semipelite. Scale card is in centimetres Gw = metagreywacke; Ps = metapsammite.

calcsilicate laminations. d) Siliceous Thompson Formation metasediments with calculicate layers and laminations from the Thompson Mine area. Cs = calcsilicate.

Manasan Formation, M1 quartzite

Paint Lake metapsammite

Setting Formation, wacke to mudstone

Anomalous Paint Lake metagreywacke

Paint Lake metapsammite

Paint Lake metagreywacke

Cs Rb Ba Th U Nb Ta K La Ce Pr Sr P Nd Zr Hf Sm Eu Ti Dy Y Yb Lu

Cs Rb Ba Th U Nb Ta K La Ce Pr Sr P Nd Zr Hf Sm Eu Ti Dy Y Yb Lu

e) Archean silicate facies iron formation with intensely folded chert layers. f) Pipe Formation P1 silicate facies iron formation at Pipe Mine. Scale card is in centimetres.

garnet, and biotite. h) Pipe Formation, P2 metapelit sillimanite-garnet-biotite gneiss from the eastern TNB. Scale card is in centimetres.

Manasan Formation, M2 semipelite

Anomalous Paint Lake metagreywacke

Paint Lake metagreywacke

Cs Rb Ba Th U Nb Ta K La Ce Pr Sr P Nd Zr Hf Sm Eu Ti Dy Y Yb Lu

Figure 2) Multi-element plots of Archean metasediments and Ospwagan Group rocks normalized to

average P2 metapelite (Zwanzig et al., 2007). Archean metasediments are compared with similar

types of lithologies from the Ospwagan Group: a) Archean metapsammite with M1 quartzite; b)

with Setting Formation quartzose wackes and mudstones. The Archean lithologies are generally

depleted in large ion lithophile elements such as Cs, Rb, Th, U, and K, and high field strength

elements such as Ta; and enriched in Sr and Eu. This suggests a more juvenile source for the

Archean metasediments such as an arc/greenstone belt. In contrast the Ospwagan Group

The Ospwagan Group geochemical data is from Zwanzig et al. (2008).

Archean metagreywacke with M2 semipelite; and c) Archean metapsammite and metagreywacke

metasediments were largely sourced from the abundant granitoids of the Archean Superior craton

i) Archean sulphide-bearing, orthopyroxene- and garnet-rich silicate facies iron formation. j) Simila orthopyroxene-bearing, sulphide- and garnet-rich silicate facies iron formation of Pipe Formation P3 member from the Owl Lake area. Gt = garnet

metapelite. The metapelite is characterized by a light gossan stain. I) Cm-scale layering of quartzo wacke and metapelite in Setting Formation from the Thompson Mine area. Pel = metapelite; Ps = metapsammite; Qw = quartzose wacke.

3) Other significant discoveries from 2009



Manitoba. Scale card is in centimetres.









Figure 3 a) Possible metavolcanics occur interbedded with metapsammite as gradationally banded mafic, intermediate, and felsic gneiss. b) A peralkaline melasyenite occurs in the SW corner of the lake and is typically metasomatized to a purplish pink to brick red colour (c). d) A swarm of pink carbonatite dykes has been extended over 21 km through the central islands of the lake. e) A possibly related series of grey silicate-carbonate dykes have also been discovered. Further information on the carbonate-rich rocks of Paint Lake can be found on the adjacent poster by Chakhmouradian et al. entitled Evidence for Carbonatite Magmatism at Paint Lake,

## 4) Economic Considerations

The similarities between the Archean supracrustal rocks and the nickel deposit-hosting Ospwagan Group supracrustal rocks could create additional difficulty for exploration in the region. The potential exists for the mistaken identification of

Ospwagan Group rocks in areas of low nickel potential. Trace element geochemistry may be useful as a tool to discriminate between the Archean supracrustals and the metasediments of the Ospwagan Group.

Alternatively, the presence of sulphide-bearing iron formation and metagreywacke in Archean supracrustal sequences could potentially make for new exploration targets. Ultramafic bodies intruding into these sulphide-bearing metasediments could utilize the available sulphur to fractionate Ni-sulphides. This is similar to the model interpreted for Ni-deposit hosting

ultramafic bodies emplaced in Ospwagan Group rocks elsewhere in the TNB.

## Acknowledgements

Thanks to J. Dutka for her assitance in the field and with sample preparation, and N. Brandson for his ever dependable logistical support. Thanks to C. Böhm, A. Chakhmouradian, T. Corkery, and J. Macek for their thoughful discussions both in and out of the

## References

uëslan, C.G. 2009: Bedrock geology of the Paint Lake area, Manitoba (parts of NTS 63O8, 9, 63P5, 12); Manitoba Innovation, Energy and Mines, Manitoba Geological Survey, Preliminary Map PMAP2009-3, scale 1:20 000.

Zwanzig, H.V., Macek, J.J., and McGregor, C.R. 2007: Lithostratigraphy and geochemistry of the high-grade metasedimentary rocks in the Thompson Nickel Belt and adjacent Kisseynew Domain, Manitoba: implications for nickel exploration; Economic Geology, v. 102, p.

Zwanzig, H.V., Macek, J.J., and McGregor, C.R. 2008: Major- and trace-element analyses of the metasedimentary and basement rocks, Thompson Nickel Belt, adjacent Superior Boundary Zone, and Kisseynew Domain; Manitoba Science, Technology, Energy, and Mines, Manitoba Geological Survey, Data Repository Item DRI2006001, Microsoft® Excel® file.

