

Figure 1. Ice-flow indicators, marine limit and limit of carbonate clasts in northeastern Manitoba. Additional data for the study area include: (a) field-based ice-flow indicators (arrows, grooves, crescentic terraces, ridges, roche moutonnée) and (b) field-based ice-flow indicators (arrows, grooves, crescentic terraces, ridges, roche moutonnée) and (c) field-based ice-flow indicators (arrows, grooves, crescentic terraces, ridges, roche moutonnée) and (d) field-based ice-flow indicators (arrows, grooves, crescentic terraces, ridges, roche moutonnée).

References: Bolton, B.S. and Clark, C.D., 1998. A highly mobile Laurentide ice sheet revealed by satellite images of glacial icebergs. Nature, v.343, p.833-837. Campbell, J.E., Trommler, M.S., McCurdy, M.W., Bohn, C.O., and Ross, M., 2012. The composition and ice-flow indicators of the Great Lakes-Caribbean Lake area (GLCC) in the northwestern Laurentide ice sheet. Geological Survey of Canada, Open File 6907, 1 CD-ROM, Manitoba Innovation, Energy and Mines. Geological Survey of Canada, Open File 6907, 1 CD-ROM, Manitoba Innovation, Energy and Mines.

Acknowledgments: This map was completed as part of a Ph.D. degree at the University of Waterloo, supervised by Dr. Ross. Funding for which was provided by the Geological Survey of Canada's (GSC) Research Affiliate Program. Field logistics and helicopter support were provided by the Manitoba Geological Survey. The mapping exercise also formed part of the GSC Geospatial for Energy and Minerals Program (GEM) Northeast Manitoba project. Natural Resources Canada, Field Geoscientist: B. Clarke, T. Stanton, R. Versey. We are grateful to the following individuals for their assistance: D. Kerr (GSC) and G. Marle (Manitoba Geological Survey) for critical review of the map.




Abstract: Northeast Manitoba is marked by glacial and geological features, with various ice-flow indicators, eskers, kames, and moraines. The map shows the distribution of these features and their relationship to the Laurentide ice sheet. The marine limit is shown as a dashed line, and the limit of carbonate clasts is shown as a solid line. The map also shows the distribution of various glacial features, including eskers, kames, and moraines.

Table with 5 columns: MAP11, MAP12, MAP13, MAP14, MAP15. It lists various map sheets and their corresponding coordinates.

National Topographic System reference and index to adjoining published maps. Cover illustration: Features and ice-flow indicators east of Kelleys Lake. Photograph by M.S. Trommler, Manitoba Geological Survey. Catalogue No. M183-141-2011E-POF. ISBN 978-1-101-92822-2.

Geological Survey of Canada, Canadian Geoscience Map 41, Manitoba Geological Survey, Geoscientific Map MAP2011-2. Stubner Lake, Manitoba, 1:50 000.



Legend and supplementary information. Includes: QUATERNARY SURFICIAL DEPOSITS (Organic, Alluvial, Lacustrine, Marine); LATE WISCONSINAN PROGLACIAL AND GLACIAL ENVIRONMENTS (Glacial, Subglacial, etc.); PRE-QUATERNARY BEDROCK (Precambrian rocks); and a list of symbols for various features like eskers, kames, and moraines. Also includes a scale bar and a north arrow.

Canadian Geoscience Maps, Cartes géoscientifiques du Canada. Includes logos for Canada and Manitoba, and contact information for the Geological Survey of Canada.

SURFICIAL GEOLOGY STUBNER LAKE Manitoba 1:50 000. Includes author information (M.S. Trommler and J.E. Campbell), publication details, and a scale bar.

Geological Survey of Canada, Canadian Geoscience Map 41, Manitoba Geological Survey, Geoscientific Map MAP2011-2. Includes contact information and a scale bar.