

GeoFile 6-2023 ReadMe

Manitoba radiocarbon ages

by M.S. Gauthier
Manitoba Geological Survey
Winnipeg, 2023



© King's Printer for Manitoba, 2023

Every possible effort is made to ensure the accuracy of the information contained in this report, but Manitoba Economic Development, Investment and Trade does not assume any liability for errors that may occur. Source references are included in the report and users should verify critical information.

Any third party digital data and software accompanying this publication are supplied on the understanding that they are for the sole use of the licensee, and will not be redistributed in any form, in whole or in part. Any references to proprietary software in the documentation and/or any use of proprietary data formats in this release do not constitute endorsement by Manitoba Economic Development, Investment and Trade of any manufacturer's product.

When using information from this publication in other publications or presentations, due acknowledgment should be given to the Manitoba Geological Survey. The following reference format is recommended:

Gauthier, M.S. 2023: Manitoba radiocarbon ages; Manitoba Economic Development, Investment and Trade, Manitoba Geological Survey, GeoFile 6-2023.

Published by:

Manitoba Economic Development, Investment and Trade
Manitoba Geological Survey
360–1395 Ellice Avenue
Winnipeg, Manitoba
R3G 3P2 Canada

Telephone: 1-800-223-5215 (General Enquiry)
204-945-6569 (Publication Sales)

Fax: 204-945-8427

Email: minesinfo@gov.mb.ca

Website: manitoba.ca/minerals

This publication supersedes Open File OF2021-1.

This publication is available to download free of charge at manitoba.ca/minerals.

This publication is available in alternate formats upon request.

Abstract

This GeoFile provides a digital dataset for radiocarbon ages obtained from samples found in Manitoba, Canada. The compilation currently has 1633 radiocarbon ages obtained for geological or archeological purposes, provided as both conventional radiocarbon ages (^{14}C years BP) and calibrated ages (cal. years BP). The dataset includes two tables with ages that have been interpreted as anomalous or maybe-anomalous. These ages should be discarded or confirmed before using them in future work. This data can be brought into GIS software, and integrated with other data, to further chronological reconstructions in Manitoba.

Résumé

Ce géodossier fournit un ensemble de données sur les âges radiocarbones obtenus à partir d'échantillons prélevés au Manitoba, au Canada. La compilation compte actuellement 1633 âges radiocarbones obtenus à des fins géologiques ou archéologiques, présentés à la fois sous forme d'âges radiocarbones conventionnels (années AP ^{14}C) et d'âges calibrés (années AP calibrées). L'ensemble de données comprend deux tables incluant des âges interprétés comme étant anomaux ou peut-être anomaux. Ces âges doivent être écartés ou confirmés avant d'être utilisés dans des travaux futurs. Ces données peuvent être incorporées au logiciel SIG et intégrées à d'autres données dans le but de compléter les reconstitutions chronologiques au Manitoba.

DIGITAL DATA

Zip file geofile6.zip contains the following folders and content:

GeoFile_6-2023_ReadMe.pdf (this file)

GeoFile_6-2023_tables.xlsx (containing Tables 4 through 9):

- Table 4: Radiocarbon ages in Manitoba.
- Table 5: Radiocarbon ages in Manitoba that need a freshwater reservoir correction.
- Table 6: 'Maybe anomalous' radiocarbon ages in Manitoba.
- Table 7: Anomalous radiocarbon ages in Manitoba.
- Table 8: References.
- Table 9: Laboratory codes.

Geofile_shps (radiocarbon shapefiles):

- Geofile_6_2023_Manitoba_radiocarbon_ages_good.shp
- Geofile_6_2023_Manitoba_radiocarbon_ages_FRE.shp
- Geofile_6_2023_Manitoba_radiocarbon_ages_maybe_anomalous.shp
- Geofile_6_2023_Manitoba_radiocarbon_ages_anomalous.shp

Introduction

Radiocarbon dating relies on the assumption that organic materials were in equilibrium with the production of ^{14}C in the atmosphere, and that the ^{14}C in the organism has decayed following the death of the organism. Because ^{14}C has a relatively short half-life (5700 ± 30 years), radiocarbon dating has a usable range of ~ 300 to $\sim 50\,000$ years (Trumbore, 2000). This GeoFile contains a compilation of radiocarbon samples analyzed at 25 laboratories between 1950 and 2022. It is both an update and revision to previous compilations (Teller, 1980; Morlan et al., 2000; Gauthier, 2021). This database will be updated annually as new data is released or reassessed. GeoFile 6 was originally released in 2023.

Methods

The Manitoba radiocarbon age database was updated and published in Gauthier (2021). That began with confirmation of data in the existing internal Manitoba Geological Survey (MGS) database used to produce Morlan et al. (2000), which was an update from Teller (1980). Resources used to update the database included internal MGS data, McNeely and Brennan (2005), Martindale et al. (2016), Dalton et al. (2020), and literature searches. Original references for each radiocarbon-age sample were verified, cited in Tables 4–7, and written in full in Table 8.

Updates since 2021

Marine age calibration

New marine–terrestrial reservoir offset (ΔR) values have been needed since the release of a new marine calibration curve (Heaton et al., 2020). Pierkowsky et al. (2022) recalculated a new ΔR for Hudson Bay should be -21 ± 72 ^{14}C years when using the Marine20 calibration curve. As such, all marine shell radiocarbon ages have been recalibrated herein.

New ages

Three samples were added from the Gillam area (UOC-16664, UOC-20687, UOC-20688), three samples from the Roseau River area (UOC-16071, UOC-16076, UOC-16077) and three samples from the Churchill River area (UOC-16584, UOC-16662, UOC-16663).

One previously-unpublished historical age from near Grunthal, discovered in MGS archives, was also added (GX-27643).

Corrections

There were a few errors in Table 4 of Open File OF2021-1 (Gauthier, 2021). As a result, the following samples have been corrected herein:

- UCIAMS-29317
- UCIAMS-88697

Amended classifications

The following classifications have been changed:

- BGS-813 to 'Maybe anomalous' due to the large error of this historical age.
- TO-4639 to 'Maybe anomalous' because this sample was not in situ and this historical age needs confirmation.
- GSC-4760 to 'Maybe anomalous' acknowledging that other regional samples are of similar age and that perhaps this is another example of interstadial sediment at surface.

Spatial characteristics

The site-location of radiocarbon samples is a mix of GPS coordinates (post-2000) and cartographic estimates (pre-2000). During the 2021 update, the elevation and coordinates of some samples were adjusted to better match the original description and/or figure given for that sample. This was achieved using a mixture of ArcMap Basemap imagery, LiDAR (where available; Government of Manitoba, 2020) and SRTM data (U.S. Geological Survey, 2014). The elevation was not updated for any 'approximate location' samples. Location adjustments were usually within 1 to 5 km. While this is not a significant distance, the corrected locations allow for better correlation to associated geomorphic landforms in a digital working environment (Gauthier et al., 2022).

Conventional normalized ages

All ages are reported as conventional radiocarbon ages (Stuiver and Polach, 1977). These ages are denoted as radiocarbon years before present (^{14}C BP), where 'present' is taken to be 1950. The error is given as a 1 sigma (σ) range for most commercial laboratories, and a 2σ range for the Geological Survey of Canada (GSC). Conventional radiocarbon ages also include a correction for isotopic fractionation ($^{13}\text{C}/^{12}\text{C}$ ratio, $\delta^{13}\text{C}$; Stuiver and Polach, 1977). This normalization is calculated using a $\delta^{13}\text{C}$ value (Stuiver and Polach, 1977) that is either measured directly by isotope ratio mass spectrometry, or provided as part of the accelerator mass spectrometry (AMS) process. Measured $\delta^{13}\text{C}$ values are provided in Tables 4–7 (Del ^{13}C). AMS-calculated values include machine fractionation and are hence not reported by the laboratories.

Estimating $\delta^{13}\text{C}$

To use historical radiocarbon ages, the user must first ensure that the data was analyzed and presented in a way that is now agreed upon by the international community (Stuiver and Polach, 1977). A problem arises when the $\delta^{13}\text{C}$ was not measured or machine-calculated, as the ages still need to be conventionally corrected (normalized) to $\delta^{13}\text{C} = -25.0\text{‰}$ (based on the Pee Dee Belemnite [PDB] standard).

Older terrestrial samples

Older terrestrial samples can be normalized using the guidance of Stuiver and Polach (1977) and Morlan (1999). Charcoal,

wood, plant macrofossils and bulk organic sediment are assumed to have been normalized to a $\delta^{13}\text{C}$ of -25.0‰ , and are hence not corrected. Peat is corrected using an assumed $\delta^{13}\text{C}$ of -27.0‰ with an error of ± 3 . Comparisons with the measured $\delta^{13}\text{C}$ values in the Manitoba dataset show that these values are generally correct (Table 1). The wood in our dataset tends to skew a bit more negative than ‘assumed’, with a mean and median $\delta^{13}\text{C}$ of -26.9‰ . There is also considerable range in the measured $\delta^{13}\text{C}$ value for bulk organic sediments, which includes everything from sandy eolian paleosols to organic lake sediment (gyttja).

Older marine samples

At GSC's radiocarbon laboratory, marine shell ages were unconventionally corrected to $\delta^{13}\text{C} = 0.0\text{‰}$ PDB instead of $\delta^{13}\text{C} = -25.0\text{‰}$ prior to 1992. This was corrected when McNeely and Brennan (2005) released corrected ages for marine shells analyzed at the GSC laboratory that had $\delta^{13}\text{C}$ measurements available. The revised shell ages are included herein.

There are 20 marine-shell ages that do not have measured $\delta^{13}\text{C}$ values (seven from the GSC lab, nine from the Brock Geosciences lab (St. Catharines, Ontario), three from Geochron Laboratories (Chelmsford, Massachusetts) and one from the University of Saskatchewan lab). Using the mean/median and standard deviation values provided by McNeely and Brennan (2005) and supported by our own measured $\delta^{13}\text{C}$ values, we have assigned ‘assumed’ $\delta^{13}\text{C}$ values according to the species of marine shell (Table 2).

There is also a marine seal bone in the database (S-521), which has been corrected using an assumed $\delta^{13}\text{C}$ value of -15‰ based on the ‘marine organisms’ estimate of Stuiver and Polach (1977).

Freshwater samples

Wherever possible, terrestrial plant macrofossils, wood and/or charcoal should be sampled for radiocarbon dating instead of aquatic plant macrofossils (cf. Marty and Myrbo, 2014), or shell-bearing organisms (ostracods, gastropods, pelecypods; cf. Philippsen, 2013). There are many times, however, where the less-ideal organic material is the only one available—the database herein contains numerous radiocarbon ages on freshwater shell-bearing organisms, as well as aquatic plant macrofossils.

Table 1: Measured $\delta^{13}\text{C}$ values for terrestrial samples in Manitoba.

	Peat	Bison bone	Human bone	Ungulate bone	Bulk organic	Charcoal	Wood
	n = 28	n = 12	n = 11	n = 4	n = 21	n = 26	n = 51
Minimum	-32.7	-25.5	-25.0	-22.1	-32.0	-28.1	-30.0
Maximum	-24.4	-15.0	-19.0	-18.1	-16.1	-20.6	-23.5
Mean	-27.5	-18.4	-21.4	-19.9	-24.8	-24.3	-26.9
Median	-27.1	-18.6	-21.0	-19.7	-25.5	-24.2	-26.9
Standard deviation	2.2	3.0	1.9	1.8	4.5	1.4	1.5

Table 2: Assumed $\delta^{13}\text{C}$ values for different taxa of marine shells, based on McNeely and Brennan (2005) and supported by measured $\delta^{13}\text{C}$ values for marine shells in Manitoba.

Species	Assumed $\delta^{13}\text{C}$ value
<i>Chlamys islandicus</i>	2.0 \pm 1.9
<i>Hiatella arctica</i>	1.2 \pm 0.7
<i>Macoma baltica</i>	-1.0 \pm 1.53
<i>Mytilus edulis</i>	0.18 \pm 1.05
Unidentified	0.85 \pm 1.27

Correcting for ^{13}C fractionation in freshwater shells is complicated since different micro-environments within the same waterbody can lead to differences in ^{13}C fractionation for different species (Fritz and Poplawski, 1974). This means that a single correction cannot be applied to all freshwater organisms, even within a single species. The variety of measured $\delta^{13}\text{C}$ values within freshwater shells in Manitoba also shows that a correction can not easily be applied (Gauthier, 2021, 2022). The same is true for ostracods (Gauthier, 2021). As such, the 25 uncorrected freshwater shell ages and two uncorrected ostracod ages herein are labelled as anomalous and should not be used.

Bones

Differences in ^{13}C fractionation between plants and grasses are further fractionated up the food chain (Morlan, 1999). Bone collagen from ungulates were corrected using an assumed $\delta^{13}\text{C}$ of $-20.0 \pm 2\text{‰}$. Ages from *Bison* sp. should be considered minimum ages, given their C_4 -plant-rich diet (Morlan, 1999). Bone collagen from *Homo sapiens* are corrected using an assumed $\delta^{13}\text{C}$ of $-19.0 \pm 2\text{‰}$.

Comparisons between the measured $\delta^{13}\text{C}$ values in the Manitoba dataset show that these values are generally correct (Table 1), and our data fits within that compiled by Morlan (1999). The human collagen in our dataset is slightly more negative than the ‘assumed’, with a mean $\delta^{13}\text{C}$ of -21.4‰ and median of -21.0‰ . The traditional diet in Manitoba was omnivorous, both for terrestrial and aquatic species (Syms, 2018). Though the data are limited, it could mean the four conventional radiocarbon ages on human bone collagen are ~30 to 40 years younger than shown (Gak-5447, S-651, S-743, S-1303). In studies where better precision is needed, it is important to ensure that the ^{13}C ratio is measured and not simply estimated. Syms (2018) suggests that this

ratio is measured for both bone collagen (protein information) and bone apatite (total diet).

Calibrated ages

To compare radiocarbon ages obtained on terrestrial, freshwater and marine organisms within Manitoba, it is necessary to calibrate the ages. It is important to note that radiocarbon ages do not directly equate with calendar years. This is because radiocarbon concentration in the atmosphere varies through time, due to changes in the production rate (de Vries, 1958). As such, calibrations use independently-dated archives such as tree rings, lacustrine and marine sediments, speleothems and corals (Heaton et al., 2020; Reimer et al., 2020). Calibrated ages are accompanied by complex, sometimes multimodal, calibrated age probability distributions that may require stratigraphic information to resolve.

All non-anomalous radiocarbon ages herein (Table 4) were calibrated using the program Calib 8.2 (Stuiver et al., 2020). Ages from terrestrial samples were calibrated using IntCal20 (Reimer et al., 2020) for the northern hemisphere. The marine mollusks were calibrated separately, to remove the reservoir effect from these shells, by using the Marine20 database (Heaton et al., 2020) with an average reservoir off-set (ΔR) of -21 ± 72 for Hudson Bay (Pieńkowski et al., 2022).

The user should note that only the highest-probability age-range is denoted herein ("Cal_BP_2 σ "), with the probability recorded next to the cal. years BP age-range in Table 4 ("Cal_BP_2 σ _probability"). Where the probability is less than 1, the user may want to refer to the probability distributions calculated within Calib 8.2 for alternate age ranges (Stuiver et al., 2020). Calibrated ages are commonly presented in the literature as median ages, with an uncertainty of 1 σ or 2 σ . The 2 σ age-ranges and the median age are included for this purpose. Ages should be presented in publications with all raw data needed for calibration (see Millard, 2014), to be updated by later researchers when new calibration curves are published. Furthermore, the convention is to round the final calculated age to the nearest 10, as the mathematical computations calculate values to a degree that is not precise in reality (Millard, 2014). When using median ages, the user must also include the 2 σ range and the probability (e.g., 8.15 ka cal. years BP, 7960–8340, 100%).

Discussion

Conventional decay-count vs. AMS methods

This GeoFile contains radiocarbon ages determined by conventional decay-count (radiometric) methods and by AMS methods. Both decay-count and AMS methods can provide comparable precision. However, decay-count requires three orders of magnitude more carbon than the AMS method. Hence, historical samples submitted for decay-count methods often used bulk sediment samples, bulk assemblages of macrofossils and/or large pieces of wood. Analysis on bulk samples can lead to

inaccuracies, as these samples may have been contaminated by reworked (older) detritus, overprinted (younger) detritus such as modern rootlets and/or contain different mixtures of materials (e.g., Bayliss and Marshall, 2019). As such, single-specimen radiocarbon samples are considered to provide more accurate results than bulk samples. At sites where organic material has been dated by both conventional and AMS methods, the AMS method is considered more accurate due to single-specimen precision.

Cautions when interpreting ages

The user is reminded that radiocarbon ages are estimates, and should always be considered with regard to other evidence from the site. For both conventional and AMS methods, trace amounts of modern carbon can generate an apparent age that is ultimately incorrect (e.g., Reyes et al., 2020). Reproducibility between (and within) laboratories is also a concern, a problem which seems to increase with older materials (Ward and Clague, 2019). In situations where a hypothesis is based on the result, duplicate radiocarbon ages should be obtained from the same material, and possibly analyzed at different labs (e.g., McMartin et al., 2019). Different pre-treatments may also help to confirm the radiocarbon age (e.g., Bajc et al., 2015). Replicate measurements on different single-specimen samples from the same context or feature can also help (Bayliss and Marshall, 2019). In all cases, other proxies (paleoenvironment, ice-flow dynamics, Heinrich events, other dating methods, cultural context, etc.) should be considered when interpreting a radiocarbon age.

Freshwater reservoir (hard-water) effect

Radiocarbon dating relies on the assumption that organic materials were in equilibrium with the production of ^{14}C in the atmosphere during their lives. Importantly, scientists have learned that organic materials are also affected by inorganic carbon within freshwater environments that overlie both carbonate rocks (Deevey et al., 1954; Andree et al., 1986; MacDonald et al., 1987), and/or lignite, coal and carbonaceous shales (Nambudiri et al., 1980). There may have been different uptake conditions within different lacustrine or fluvial bodies, and within different time periods (Shotton, 1972). Contamination by old-carbon is termed the hard-water effect, and is important in Manitoba because most glacial and postglacial sediments in Manitoba are calcareous (Manitoba Natural Resources and Northern Development, 2022). The hard-water effect is also important when interpreting 'terrestrial' radiocarbon ages obtained from bones of species (birds, humans, canines, bear, wolf, etc.) that may have eaten a marine and/or freshwater diet (Syms, 2018). In Manitoba, the freshwater diet effect ($n = 4$) varies from 220 to 370 ^{14}C years on specimens that are between 440 and 4400 ^{14}C years (Gauthier, 2022). For newer analyses, $^{15}\text{N}/^{14}\text{N}$ ratios should be analyzed to help determine if fish or other aquatic species were part of the diet—if they were, a freshwater (hard-water) correction is needed (Syms, 2018). The column 'Del ^{15}N ' has been added, and it is the intention of the MGS to collect this ratio on all bones and antlers moving forward.

A hard-water correction is typically calculated by comparing the ages of terrestrial and freshwater material collected from the same horizon at the same site. For Manitoba, the hardwater effect ($n = 12$) is temporally variable, with a median of 230 ^{14}C years and values up to at least 880 ^{14}C years (Gauthier, 2022). Eleven of those paired terrestrial-freshwater ages are less than 5000 ^{14}C years old, meaning an accurate hardwater correction for older material is very difficult to determine. To acknowledge the lack of regionally- and temporally-specific data, no hard-water corrections have been made to the data herein. Instead, the notation “FRE?” is added to the conventional and calibrated ages in Table 5 (123 ages), acknowledging the need for future correction as the user finds appropriate. Potential bone samples affected by a freshwater diet (e.g., *Homo sapiens*), and without a paired terrestrial sample, are also included in this table.

Marine reservoir effect

The concentration of carbon isotopes differs between the ocean and the atmosphere, with a ‘reservoir’ in the global oceans that results from trapped old carbon (Arnold, 1957; de Vries, 1958). This reservoir effect results in an apparent radiocarbon age of a marine sample that is different from the true age (when carbon exchange between the organism and the atmosphere would be equal). As such, measured radiocarbon ages from marine samples can’t be directly compared to measured radiocarbon ages from terrestrial samples. Complicating matters, the level of depletion varies due to spatio-temporal differences in the ocean and the atmosphere (Heaton et al., 2020). To account for this, a marine radiocarbon reservoir age is applied to marine organisms used in radiocarbon dating (Stuiver et al., 1986; McNeely et al., 2006; Coulthard et al., 2010). The marine radiocarbon reservoir is calculated using the difference between the global mean ocean reservoir correction and the regional reservoir age, termed the regional reservoir offset ($\Delta R(\theta)$; Heaton et al., 2020). $\Delta R(\theta)$ will remain approximately constant through time, assuming that the regional oceanographic characteristics remain (Stuiver et al., 1986; Stuiver and Braziunas, 1993). Herein, we calibrate the marine radiocarbon ages using the mollusc marine regional reservoir offset $\Delta R(\theta) = -21 \pm 72$ years, for Hudson Bay (Pieńkowski et al., 2022). The regional mollusc marine radiocarbon reservoir (R_r) was calculated using twelve live-collected shells sampled between 1920 and 1954 (pre-bomb).

It should be noted that most Canadian marine radiocarbon reservoir studies involve deposit-feeding molluscs. Recent studies on marine walrus, beluga whale and bowhead whale bones/teeth have determined different reservoir offsets than that of molluscs (Dyke et al., 2019; Pieńkowski et al., 2022), which should caution the user against using mollusc-based corrections for marine mammals. The database herein has one single marine mammal age, which for lack of a better correction, has been tentatively calibrated using the mollusc marine reservoir age (S-521, Rutherford et al., 1973). This calibration is likely wrong.

Marine hard-water effect

Marine materials are affected by inorganic carbon within environments that overlie carbonate-bearing or calcareous rocks or sediment. Open marine water contains dissolved inorganic carbon, which generally masks the hard-water effect. The hard-water effect may be a concern in areas with restricted water circulation, areas of considerable mixing between freshwater and marine waters, overlying highly calcareous substrate, and in areas with high abundance of terrestrial organic matter (Douka et al., 2010). Radiocarbon dating of filter-feeding molluscs will result in avoidance of the hard-water effect relative to deposit-feeding molluscs (England et al., 2013). The database herein contains radiocarbon ages for eight different shell taxa, broken down in Table 3. A ninth taxon, *Portlandia arctica* can be found within the anomalous table, given its proven hard-water effects (England et al., 2013). The twelve *Macoma* sp. radiocarbon ages herein are tagged as ‘Maybe anomalous’ (Table 6), because they are deposit feeders and may need a hard-water correction. Unidentified shells are also classified as ‘maybe anomalous’, given the difficulty of determining feeding behaviour on these ages. Only one site in Manitoba contains radiocarbon ages on bulk samples of both *Hiattella arctica* and *Macoma baltica*. As the *Hiattella arctica* samples (GSC-3367, BGS-791; Nielsen et al., 1986) were collected ~4 m lower than the *Macoma baltica* sample (BGS-797; Morlan et al., 2000), it is impossible to assess a potential hard-water effect—except to note that the upper shells are younger, as expected.

There are very few early Holocene paired terrestrial-marine samples within Hudson Bay, with which to assess a hard-water effect. On Southampton Island, early–mid-Holocene paired terrestrial-marine ages provided a regional marine reservoir age of -630 ± 45 ^{14}C years (Ross et al., 2012) and a reservoir offset (ΔR) age of 263 ± 48 years. That is higher than the ΔR age of 175 ± 89 years for the Foxe basin proposed by Pieńkowski et al. (2022), and suggests the modern correction may need a hard-water

Table 3: Marine shell taxa within the Manitoba radiocarbon database, and their feeding behaviours (after McNeely et al., 2006).

Species	Number of samples	Habitat ¹	Feeding ²
<i>Astarte borealis</i>	1	Infaunal	Suspension
<i>Chlamys islandicus</i>	3	Epifaunal	Suspension
<i>Clinocardium ciliatum</i>	1	Infaunal	Suspension
<i>Hiattella arctica</i>	40	Epifaunal	Suspension
<i>Macoma baltica</i>	2	Infaunal	Deposit
<i>Macoma calcarea</i>	10	Infaunal	Deposit
<i>Mya truncata</i>	4	Epifaunal	Suspension
<i>Mytilus edulis</i>	9	Epifaunal	Suspension
Unidentified	21	?	?

¹ Epifauna are organisms that live upon the surface of sediments. Infauna are organisms that live within sediments.
² Suspension feeders (filter feeders) are animals that feed by straining suspended matter from the water. Deposit feeders are animals that feed by obtaining food particles in the sediment.

correction. Contrastingly, on Baffin Island in the eastern Foxe basin, Vickers et al. (2010) calculated a mean ΔR age of 615 ± 20 years for early–mid Holocene terrestrial-marine pairs. Ross et al. (2012) suggested the difference may be due to underlying geology (the hard-water effect), as northern Southampton Island is granitoid while the Baffin Island sites are calcareous. As such, more local data is needed to ascertain the validity of the corrections applied herein over both time and space.

Infinite and near-infinite ages

Very old samples have such low concentrations of ^{14}C that they cannot be distinguished reliably from the background radiation. Different laboratories set different ‘ages’ as the upper limit, which can vary based on the weight of the material submitted (Pigati et al., 2007); these ages are usually reported as $> xx$ years BP. Regardless of the reported number, the age of these samples was not determined using radiocarbon methods and is interpreted as non-finite (greater than).

As laboratory methods improve over time, the upper boundary of infinite ages has increased. The user is reminded that small amounts of contamination by younger carbon will have large effects near the upper bounds of radiocarbon limits. Contamination can be introduced during burial (diagenetic or modern root-lets), surface weathering, sampling and/or laboratory processing (Pigati et al., 2007). Hence, a lab-accurate age of 48 ka ^{14}C BP may not be a ‘true’ accurate age for that organism. Contamination is more likely for peat (modern roots) or carbonate shells (recrystallization; Douka et al., 2010), than for wood. Recommendations on the ‘validity’ of near-infinite ages range from ~ 40 ka (Walker, 2005) to ~ 35 – 40 ka (Miller and Andrews, 2019)—though the latter paper lacks a source for such a recommendation. In every case, all available geological and site-specific evidence should be considered when interpreting age results, and other proxies (paleoenvironment, ice-flow dynamics, Heinrich events, other dating methods, etc.) should be considered when interpreting the radiocarbon age.

Anomalous and maybe-anomalous ages

A number of radiocarbon ages have been identified as anomalous, either by the author during the course of interpretation, or if the material was bulk organic sediment, lake sediment, or organic mud. The latter are tagged as anomalous given the problems with conventional dating of large mixed samples that may contain older transported detritus and/or younger intruding detritus (e.g., Clayton and Moran, 1982; Grimm et al., 2009; Bayliss and Marshall, 2019). Bulk lake sediment (gyttja) samples also contain nonorganic detritus that contributes to a hard-water effect (see below), which is difficult to correct for given the potential for contamination and re-working. A recent compilation paper by Young et al. (2021) has chosen to reject all bulk lake/soil samples during their analysis—and we advise the same. These rejected samples are retained in the database, and clearly marked ‘yes’ under the column ‘ANOMALOUS’ (263 samples, Table 7). The notation “A” is added to the conventional age, to

further remind the user that the age is likely not valid. The specific reason why each sample was determined to be anomalous is written under column ‘ANOMALOUS_WHY’.

Additional samples are tagged as ‘maybe’ anomalous. This includes stratigraphically inversed (e.g., older above younger) ages and ages near the upper boundaries for radiocarbon dating (>40 ka). Near-finite ages should be confirmed through stratigraphy (including till composition) and re-dating, since they are near the boundary between interstadial marine isotope stage (MIS) 3 (29–57 cal. ka BP; Lisiecki and Raymo, 2005), and MIS 5 (infinite). The notation “M” is added to the conventional age, to further remind the user that the age may not be valid (123 samples, Table 6). The specific reason why each sample was determined to be maybe anomalous is written under column ‘ANOMALOUS_WHY’.

Macrofossil reports

A number of samples were submitted to A. Telka for macrofossil identification. Most of these samples are associated with a macrofossil report, which includes information on different types of plant and animal macrofossils contained within the submitted sediment. These reports are identified with a number under the column “REPORT” and the actual report information is embedded within the “COMMENTS” column of Tables 4–7.

References

- Andree, M., Oeschger, H., Siegenthaler, U., Riesen, T., Moell, M., Ammann, B. and Tobolski, K. 1986: ^{14}C dating of plant macrofossils in lake sediment; *Radiocarbon*, v. 28, no. 2A, p. 411–416.
- Arnold, J.R. 1957: The distribution of carbon-14 in nature; *Tellus*, v. 9, no. 1, p. 28–32.
- Bajc, A.F., Karrow, P.F., Yansa, C.H., Curry, B.B., Nekola, J.C., Seymour, K.L. and Mackie, G.L. 2015: Geology and paleoecology of a Middle Wisconsin fossil occurrence in Zorra Township, southwestern Ontario, Canada; *Canadian Journal of Earth Sciences*, v. 52, no. 6, p. 386–404, URL <<https://doi.org/10.1139/cjes-2015-0005>>.
- Bayliss, A. and Marshall, P. 2019: Confessions of a serial polygamist: the reality of radiocarbon reproducibility in archaeological samples; *Radiocarbon*, v. 61, no. 5, p. 1143–1158, URL <<https://doi.org/10.1017/RDC.2019.55>>.
- Clayton, L. and Moran, S.R. 1982: Chronology of late Wisconsinan glaciations in middle North America; *Quaternary Science Reviews*, v. 1, p. 55–82.
- Coulthard, R.D., Furze, M.F.A., Pieńkowski, A.J., Nixon, F.C. and England, J.H. 2010: New marine ΔR values for Arctic Canada; *Quaternary Geochronology*, v. 5, no. 4, p. 419–434.
- Dalton, A.S., Margold, M., Stokes, C.R., Tarasov, L., Dyke, A.S., Adams, R.S., Allard, S., Arends, H., Atkinson, N., Attig, J.W., Barnett, P.J., Barnett, R.L., Batterson, M., Bernatchez, P., Borns, H.W., Breckenridge, A., Briner, J.P., Brouard, E., Campbell, J.E., Carlson, A.E. et al. 2020: An updated radiocarbon-based ice margin chronology for the last deglaciation of the North American Ice Sheet Complex; *Quaternary Science Reviews*, v. 234, art. 106223, URL <<https://doi.org/10.1016/j.quascirev.2020.106223>>.
- de Vries, H. 1958: Variations in concentration of radiocarbon with time and location on earth; *Koninklijke Nederlandse Akademie van Wetenschappen, Proc. Ser. B*, v. 61, no. 2, p. 94–102.

- Deevey, E.S., Jr., Gross, M.S., Hutchinson, G.E. and Kraybill, H.L. 1954: The natural C^{14} contents of materials from hard-water lakes; *PNAS*, v. 40, no. 5, p. 285–288, URL <<https://doi.org/10.1073/pnas.40.5.285>>.
- Douka, K., Higham, T.F.G. and Hedges, R.E.M. 2010: Radiocarbon dating of shell carbonates: old problems and new solutions; *Munibe Suplemento*, v. 31, p. 18–27.
- Dyke, A.S., Saville, J.M., Szpak, P., Southon, J.R., Howse, L., Desrosiers, P.M. and Kotar, K. 2019: An assessment of marine reservoir corrections for radiocarbon dates on walrus from the Foxe Basin region of Arctic Canada; *Radiocarbon*, v. 61, no. 1, p. 67–81, URL <<https://doi.org/10.1017/RDC.2018.50>>.
- England, J., Dyke, A.S., Coulthard, R.D., McNeely, R. and Aitken, A. 2012: The exaggerated radiocarbon age of deposit-feeding molluscs in calcareous environments; *Boreas*, v. 42, no. 2, p. 362–373, URL <<https://doi.org/10.1111/j.1502-3885.2012.00256.x>>.
- Fritz, P. and Poplawski, S. 1974: ^{18}O and ^{13}C in the shells of freshwater molluscs and their environments; *Earth and Planetary Science Letters*, v. 24, no. 1, p. 91–98.
- Gauthier, M.S. 2021: Manitoba radiocarbon ages: update; Manitoba Agriculture and Resource Development, Manitoba Geological Survey, Open File OF2021-1, 7 p. plus 2 appendices.
- Gauthier, M.S. 2022: Using radiocarbon ages on organics affected by freshwater—a geologic and archaeologic update on the freshwater reservoir ages and freshwater diet effect in Manitoba, Canada; *Radiocarbon*, v. 64, no. 2, p. 253–264, URL <<https://doi.org/10.1017/RDC.2022.30>>.
- Gauthier, M.S., Santucci, A. and Keller, G.R. 2022: Digital compilation of surficial point and line features for Manitoba, including ice-flow data; Manitoba Natural Resources and Northern Development, Manitoba Geological Survey, GeoFile 1-2022.
- Government of Manitoba 2020: Southern Manitoba LIDAR data; URL <https://mli2.gov.mb.ca/dems/index_external_lidar.html> [January 2020].
- Grimm, E.C., Maher, L.J., Jr. and Nelson, D.M. 2009: The magnitude of error in conventional bulk-sediment radiocarbon dates from central North America; *Quaternary Research*, v. 72, no. 2, p. 301–308.
- Heaton, T.J., Kohler, P., Butzin, M., Bard, E., Reimer, R.W., Austin, W. E.N., Bronk Ramsey, C., Grootes, P.M., Hughen, K.A., Kromer, B., Reimer, P.J., Adkins, J., Burke, A., Cook, M.S., Olsen, J. and Skinner, L.C. 2020: Marine20—the marine radiocarbon age calibration curve (0–55,000 cal BP); *Radiocarbon*, v. 62, no. 4, p. 779–820, URL <<https://doi.org/10.1017/RDC.2020.68>>.
- Lisiecki, L.E. and Raymo, M.E. 2005: A Pliocene-Pleistocene stack of 57 globally distributed benthic $\delta^{18}O$ records; *Paleoceanography and Paleoclimatology*, v. 20, no. 1, art. PA1003.
- MacDonald, G.M., Beukens, R.P., Kieser, W.E. and Vitt, D.H. 1987: Comparative radiocarbon dating of terrestrial plant macrofossils and aquatic moss from the “ice-free corridor” of western Canada; *Geology*, v. 15, no. 9, p. 837–840, URL <[https://doi.org/10.1130/0091-7613\(1987\)15<837:CRDOTP>2.0.CO;2](https://doi.org/10.1130/0091-7613(1987)15<837:CRDOTP>2.0.CO;2)>.
- Manitoba Natural Resources and Northern Development 2022: Manitoba carbonate dispersal analyses in till; Manitoba Natural Resources and Northern Development, Manitoba Geological Survey, URL <<http://manitoba.ca/iem/geo/surficial/carbonate.html>> [July 2022].
- Martindale, A., Morlan, R., Betts, M., Blake, M., Gajewski, K., Chaput, M., Mason, A. and Vermeersch, P. 2016: Canadian archaeological radiocarbon database (CARD 2.1); URL <<https://www.canadianarchaeology.ca>> [December 2020].
- Marty, J. and Myrbo, A. 2014: Radiocarbon dating suitability of aquatic plant macrofossils; *Journal of Paleolimnology*, v. 52, p. 435–443.
- McMartin, I., Campbell, J.E. and Dredge, L.A. 2019: Middle Wisconsinan marine shells near Repulse Bay, Nunavut, Canada: implications for marine isotope stage 3 ice-free conditions and Laurentide Ice Sheet dynamics in north-west Hudson Bay; *Journal of Quaternary Science*, v. 34, no. 1, p. 64–75, URL <<https://doi.org/10.1002/jqs.3081>>.
- McNeely, R. and Brennan, J. 2005: Geological Survey of Canada revised shell dates; Geological Survey of Canada, Open File 5019, 530 p.
- McNeely, R., Dyke, A.S. and Southon, J.R. 2006: Canadian marine reservoir ages, preliminary data assessment; Geological Survey of Canada, Open File 5049, 3 p.
- Millard, A.R. 2014: Conventions for reporting radiocarbon determinations; *Radiocarbon*, v. 56, no. 2, p. 555–559, URL <<https://doi.org/10.2458/56.17455>>.
- Miller, G.H. and Andrews, J.T. 2019: Hudson Bay was not deglaciated during MIS-3; *Quaternary Science Reviews*, v. 225, no. 1, art. 105944, URL <<https://doi.org/10.1016/j.quascirev.2019.105944>>.
- Morlan, R. 1999: Canadian archaeological radiocarbon database: establishing conventional ages; *Canadian Journal of Archaeology*, v. 23, no. 1/2, p. 3–10, URL <<https://www.jstor.org/stable/41103370>> [June 2023].
- Morlan, R.E., McNeely, R. and Nielsen, E. 2000: Manitoba radiocarbon dates; Manitoba Industry, Trade and Mines, Manitoba Geological Survey, Open File Report OF2000-1, 198 p.
- Nambudiri, E.M.V., Teller, J.T. and Last, W.M. 1980: Pre-Quaternary microfossils—a guide to errors in radiocarbon dating; *Geology*, v. 8, no. 3, p. 123–126, URL <[https://doi.org/10.1130/0091-7613\(1980\)8<123:PMGTET>2.0.CO;2](https://doi.org/10.1130/0091-7613(1980)8<123:PMGTET>2.0.CO;2)>.
- Nielsen, E., Morgan, A.V., Morgan, A., Mott, R.J., Rutter, N.W. and Causse, C. 1986: Stratigraphy, paleoecology and glacial history of the Gillam area, Manitoba; *Canadian Journal of Earth Sciences*, v. 23, no. 11, p. 1641–1661.
- Philippson, B. 2013: The freshwater reservoir effect in radiocarbon dating; *Heritage Science*, v. 1, no. 24, 19 p., URL <<https://heritagesciencejournal.springeropen.com/articles/10.1186/2050-7445-1-24>> [June 2023].
- Pieńkowski, A.J., Coulthard, R.D. and Furze, M.F. A. 2022: Revised marine reservoir offset (ΔR) values for molluscs and marine mammals from Arctic North America; *Boreas*, v. 52, no. 2, p. 145–167, URL <<https://doi.org/10.1111/bor.12606>>.
- Pigati, J.S., Quade, J., Wilson, J., Jull, A.J.T. and Lifton, N.A. 2007: Development of low-background vacuum extraction and graphitization systems for ^{14}C dating of old (40–60ka) samples; *Quaternary International*, v. 166, no. 1, p. 4–14.
- Reimer, P.J., Austin, W.E.N., Bard, E., Bayliss, A., Blackwell, P.G., Bronk Ramsey, C., Butzin, M., Cheng, H., Edwards, R.L., Friedrich, M., Grootes, P.M., Guilderson, T.P., Hajdas, I., Heaton, T.J., Hogg, A.G., Hughen, K.A., Kromer, B., Manning, S., Muscheler, R., Palmer, J.G. et al. 2020: The IntCal20 northern hemisphere radiocarbon age calibration curve (0–55 cal kBP); *Radiocarbon*, v. 62, no. 4, p. 725–757, URL <<https://doi.org/10.1017/RDC.2020.41>>.
- Reyes, A.V., Dillman, T., Kennedy, K., Froese, D., Beaudoin, A.B. and Paulen, R.C. 2020: Legacy radiocarbon ages and the MIS 3 dating game: a cautionary tale from re-dating of pre-LGM sites in western Canada; Geological Society of America, GSA 2020 Connects Online virtual event, October 26–30, 2020, Abstracts with Programs, v. 52, no. 6, p. 1, paper no. 56-3, URL <<https://doi.org/10.1130/abs/2020AM-360064>>.
- Ross, M., Utting, D.J., Lajeunesse, P. and Kosar, K.G.A. 2012: Early Holocene deglaciation of northern Hudson Bay and Foxe Channel constrained by new radiocarbon ages and marine reservoir correction; *Quaternary Research*, v. 78, no. 1, p. 82–94.

- Rutherford, A.A., Wittenberg, J. and McCallum, K.J. 1973: University of Saskatchewan radiocarbon dates VI; *Radiocarbon*, v. 15, no. 1, p. 193–211.
- Shotton, F.W. 1972: An example of hard-water error in radiocarbon dating of vegetable matter; *Nature*, v. 240, p. 460–461, URL <<https://doi.org/10.1038/240460a0>>.
- Stuiver, M. and Braziunas, T.F. 1993: Modeling atmospheric ^{14}C influences and ^{14}C ages of marine samples to 10,000 BC; *Radiocarbon*, v. 35, no. 1, p. 137–189, URL <<https://doi.org/10.1017/S0033822200013874>>.
- Stuiver, M. and Polach, H.A. 1977: Discussion reporting of ^{14}C data; *Radiocarbon*, v. 19, no. 3, p. 355–363.
- Stuiver, M., Pearson, G.W. and Braziunas, T. 1986: Radiocarbon age calibration of marine samples back to 9000 cal yr BP; *Radiocarbon*, v. 28, no. 2B, p. 980–1021.
- Stuiver, M., Reimer, P.J. and Reimer, R. 2020: CALIB 8.2 [WWW program]; URL <<http://calib.org>> [December 2020].
- Syms, E.L. 2018: Discovering the fresh water reservoir effect in the northern boreal forests: refining the temporal framework; *Ontario Archaeology*, no. 98, p. 48–65.
- Teller, J.T. 1980: Radiocarbon dates in Manitoba; Manitoba Department of Energy and Mines, Mineral Resources Division, Geological Report GR80-4, 61 p.
- Trumbore, S.E. 2000: Radiocarbon geochronology; in *Quaternary Geochronology: methods and applications*, J.S. Noller, J.M. Sowers and W.R. Lettis (ed.), American Geophysical Union, Washington, D.C., p. 41–60.
- U.S. Geological Survey 2014: USGS EROS archive - digital elevation - Shuttle Radar Topography Mission (SRTM) 1 arc-second global; U.S. Geological Survey, 30 m cell, zipped hgt format, URL <<https://doi.org/10.5066/F7PR7TFT>>.
- Vickers, K.J., Ward, B.C., Utting, D.J. and Telka, A.M. 2010: Deglacial reservoir age and implications, Foxe Peninsula, Baffin Island; *Journal of Quaternary Science*, v. 25, p. 1388–1346, URL <<https://doi.org/10.1002/jqs.1419>>.
- Walker, M. 2005: *Quaternary dating methods*; John Wiley & Sons, Ltd., Chichester, West Sussex, United Kingdom, 304 p.
- Ward, B.C. and Clague, J. 2019: A blind comparison of radiocarbon labs; European Geosciences Union, EGU General Assembly 2019, April 7–12, 2019, Vienna, Austria, Geophysical Research Abstracts, v. 21, EGU2019-12003.
- Young, J.M., Reyes, A.V. and Froese, D.G. 2021: Assessing the ages of the Moorhead and Emerson phases of glacial Lake Agassiz and their temporal connection to the Younger Dryas cold reversal; *Quaternary Science Reviews*, v. 251, art. 106714, URL <<https://doi.org/10.1016/j.quascirev.2020.106714>>.

Abstract

This GeoFile provides a digital dataset for radiocarbon ages obtained from samples found in Manitoba, Canada. The compilation currently has 1633 radiocarbon ages obtained for geological or archeological purposes, provided as both conventional radiocarbon ages (^{14}C years BP) and calibrated ages (cal. years BP). The dataset includes two tables with ages that have been interpreted as anomalous or maybe-anomalous. These ages should be discarded or confirmed before using them in future work. This data can be brought into GIS software, and integrated with other data, to further chronological reconstructions in Manitoba.

Résumé

Ce géodossier fournit un ensemble de données sur les âges radiocarbones obtenus à partir d'échantillons prélevés au Manitoba, au Canada. La compilation compte actuellement 1633 âges radiocarbones obtenus à des fins géologiques ou archéologiques, présentés à la fois sous forme d'âges radiocarbones conventionnels (années AP ^{14}C) et d'âges calibrés (années AP calibrées). L'ensemble de données comprend deux tables incluant des âges interprétés comme étant anomaux ou peut-être anomaux. Ces âges doivent être écartés ou confirmés avant d'être utilisés dans des travaux futurs. Ces données peuvent être incorporées au logiciel SIG et intégrées à d'autres données dans le but de compléter les reconstitutions chronologiques au Manitoba.

DIGITAL DATA

Zip file geofile6.zip contains the following folders and content:

GeoFile_6-2023_ReadMe.pdf (this file)

GeoFile_6-2023_tables.xlsx (containing Tables 4 through 9):

- Table 4: Radiocarbon ages in Manitoba.
- Table 5: Radiocarbon ages in Manitoba that need a freshwater reservoir correction.
- Table 6: 'Maybe anomalous' radiocarbon ages in Manitoba.
- Table 7: Anomalous radiocarbon ages in Manitoba.
- Table 8: References.
- Table 9: Laboratory codes.

Geofile_shps (radiocarbon shapefiles):

- Geofile_6_2023_Manitoba_radiocarbon_ages_good.shp
- Geofile_6_2023_Manitoba_radiocarbon_ages_FRE.shp
- Geofile_6_2023_Manitoba_radiocarbon_ages_maybe_anomalous.shp
- Geofile_6_2023_Manitoba_radiocarbon_ages_anomalous.shp

Introduction

Radiocarbon dating relies on the assumption that organic materials were in equilibrium with the production of ^{14}C in the atmosphere, and that the ^{14}C in the organism has decayed following the death of the organism. Because ^{14}C has a relatively short half-life (5700 ± 30 years), radiocarbon dating has a usable range of ~ 300 to $\sim 50\,000$ years (Trumbore, 2000). This GeoFile contains a compilation of radiocarbon samples analyzed at 25 laboratories between 1950 and 2022. It is both an update and revision to previous compilations (Teller, 1980; Morlan et al., 2000; Gauthier, 2021). This database will be updated annually as new data is released or reassessed. GeoFile 6 was originally released in 2023.

Methods

The Manitoba radiocarbon age database was updated and published in Gauthier (2021). That began with confirmation of data in the existing internal Manitoba Geological Survey (MGS) database used to produce Morlan et al. (2000), which was an update from Teller (1980). Resources used to update the database included internal MGS data, McNeely and Brennan (2005), Martindale et al. (2016), Dalton et al. (2020), and literature searches. Original references for each radiocarbon-age sample were verified, cited in Tables 4–7, and written in full in Table 8.

Updates since 2021

Marine age calibration

New marine–terrestrial reservoir offset (ΔR) values have been needed since the release of a new marine calibration curve (Heaton et al., 2020). Pierńkowski et al. (2022) recalculated a new ΔR for Hudson Bay should be -21 ± 72 ^{14}C years when using the Marine20 calibration curve. As such, all marine shell radiocarbon ages have been recalibrated herein.

New ages

Three samples were added from the Gillam area (UOC-16664, UOC-20687, UOC-20688), three samples from the Roseau River area (UOC-16071, UOC-16076, UOC-16077) and three samples from the Churchill River area (UOC-16584, UOC-16662, UOC-16663).

One previously-unpublished historical age from near Grunthal, discovered in MGS archives, was also added (GX-27643).

Corrections

There were a few errors in Table 4 of Open File OF2021-1 (Gauthier, 2021). As a result, the following samples have been corrected herein:

- UCIAMS-29317
- UCIAMS-88697

Amended classifications

The following classifications have been changed:

- BGS-813 to 'Maybe anomalous' due to the large error of this historical age.
- TO-4639 to 'Maybe anomalous' because this sample was not in situ and this historical age needs confirmation.
- GSC-4760 to 'Maybe anomalous' acknowledging that other regional samples are of similar age and that perhaps this is another example of interstadial sediment at surface.

Spatial characteristics

The site-location of radiocarbon samples is a mix of GPS coordinates (post-2000) and cartographic estimates (pre-2000). During the 2021 update, the elevation and coordinates of some samples were adjusted to better match the original description and/or figure given for that sample. This was achieved using a mixture of ArcMap Basemap imagery, LiDAR (where available; Government of Manitoba, 2020) and SRTM data (U.S. Geological Survey, 2014). The elevation was not updated for any 'approximate location' samples. Location adjustments were usually within 1 to 5 km. While this is not a significant distance, the corrected locations allow for better correlation to associated geomorphic landforms in a digital working environment (Gauthier et al., 2022).

Conventional normalized ages

All ages are reported as conventional radiocarbon ages (Stuiver and Polach, 1977). These ages are denoted as radiocarbon years before present (^{14}C BP), where 'present' is taken to be 1950. The error is given as a 1 sigma (σ) range for most commercial laboratories, and a 2σ range for the Geological Survey of Canada (GSC). Conventional radiocarbon ages also include a correction for isotopic fractionation ($^{13}\text{C}/^{12}\text{C}$ ratio, $\delta^{13}\text{C}$; Stuiver and Polach, 1977). This normalization is calculated using a $\delta^{13}\text{C}$ value (Stuiver and Polach, 1977) that is either measured directly by isotope ratio mass spectrometry, or provided as part of the accelerator mass spectrometry (AMS) process. Measured $\delta^{13}\text{C}$ values are provided in Tables 4–7 (Del ^{13}C). AMS-calculated values include machine fractionation and are hence not reported by the laboratories.

Estimating $\delta^{13}\text{C}$

To use historical radiocarbon ages, the user must first ensure that the data was analyzed and presented in a way that is now agreed upon by the international community (Stuiver and Polach, 1977). A problem arises when the $\delta^{13}\text{C}$ was not measured or machine-calculated, as the ages still need to be conventionally corrected (normalized) to $\delta^{13}\text{C} = -25.0\text{‰}$ (based on the Pee Dee Belemnite [PDB] standard).

Older terrestrial samples

Older terrestrial samples can be normalized using the guidance of Stuiver and Polach (1977) and Morlan (1999). Charcoal,

wood, plant macrofossils and bulk organic sediment are assumed to have been normalized to a $\delta^{13}\text{C}$ of -25.0‰ , and are hence not corrected. Peat is corrected using an assumed $\delta^{13}\text{C}$ of -27.0‰ with an error of ± 3 . Comparisons with the measured $\delta^{13}\text{C}$ values in the Manitoba dataset show that these values are generally correct (Table 1). The wood in our dataset tends to skew a bit more negative than ‘assumed’, with a mean and median $\delta^{13}\text{C}$ of -26.9‰ . There is also considerable range in the measured $\delta^{13}\text{C}$ value for bulk organic sediments, which includes everything from sandy eolian paleosols to organic lake sediment (gyttja).

Older marine samples

At GSC's radiocarbon laboratory, marine shell ages were unconventionally corrected to $\delta^{13}\text{C} = 0.0\text{‰}$ PDB instead of $\delta^{13}\text{C} = -25.0\text{‰}$ prior to 1992. This was corrected when McNeely and Brennan (2005) released corrected ages for marine shells analyzed at the GSC laboratory that had $\delta^{13}\text{C}$ measurements available. The revised shell ages are included herein.

There are 20 marine-shell ages that do not have measured $\delta^{13}\text{C}$ values (seven from the GSC lab, nine from the Brock Geosciences lab (St. Catharines, Ontario), three from Geochron Laboratories (Chelmsford, Massachusetts) and one from the University of Saskatchewan lab). Using the mean/median and standard deviation values provided by McNeely and Brennan (2005) and supported by our own measured $\delta^{13}\text{C}$ values, we have assigned ‘assumed’ $\delta^{13}\text{C}$ values according to the species of marine shell (Table 2).

There is also a marine seal bone in the database (S-521), which has been corrected using an assumed $\delta^{13}\text{C}$ value of -15‰ based on the ‘marine organisms’ estimate of Stuiver and Polach (1977).

Freshwater samples

Wherever possible, terrestrial plant macrofossils, wood and/or charcoal should be sampled for radiocarbon dating instead of aquatic plant macrofossils (cf. Marty and Myrbo, 2014), or shell-bearing organisms (ostracods, gastropods, pelecypods; cf. Philippsen, 2013). There are many times, however, where the less-ideal organic material is the only one available—the database herein contains numerous radiocarbon ages on freshwater shell-bearing organisms, as well as aquatic plant macrofossils.

Table 1: Measured $\delta^{13}\text{C}$ values for terrestrial samples in Manitoba.

	Peat	Bison bone	Human bone	Ungulate bone	Bulk organic	Charcoal	Wood
	n = 28	n = 12	n = 11	n = 4	n = 21	n = 26	n = 51
Minimum	-32.7	-25.5	-25.0	-22.1	-32.0	-28.1	-30.0
Maximum	-24.4	-15.0	-19.0	-18.1	-16.1	-20.6	-23.5
Mean	-27.5	-18.4	-21.4	-19.9	-24.8	-24.3	-26.9
Median	-27.1	-18.6	-21.0	-19.7	-25.5	-24.2	-26.9
Standard deviation	2.2	3.0	1.9	1.8	4.5	1.4	1.5

Table 2: Assumed $\delta^{13}\text{C}$ values for different taxa of marine shells, based on McNeely and Brennan (2005) and supported by measured $\delta^{13}\text{C}$ values for marine shells in Manitoba.

Species	Assumed $\delta^{13}\text{C}$ value
<i>Chlamys islandicus</i>	2.0 \pm 1.9
<i>Hiatella arctica</i>	1.2 \pm 0.7
<i>Macoma baltica</i>	-1.0 \pm 1.53
<i>Mytilus edulis</i>	0.18 \pm 1.05
Unidentified	0.85 \pm 1.27

Correcting for ^{13}C fractionation in freshwater shells is complicated since different micro-environments within the same waterbody can lead to differences in ^{13}C fractionation for different species (Fritz and Poplawski, 1974). This means that a single correction cannot be applied to all freshwater organisms, even within a single species. The variety of measured $\delta^{13}\text{C}$ values within freshwater shells in Manitoba also shows that a correction can not easily be applied (Gauthier, 2021, 2022). The same is true for ostracods (Gauthier, 2021). As such, the 25 uncorrected freshwater shell ages and two uncorrected ostracod ages herein are labelled as anomalous and should not be used.

Bones

Differences in ^{13}C fractionation between plants and grasses are further fractionated up the food chain (Morlan, 1999). Bone collagen from ungulates were corrected using an assumed $\delta^{13}\text{C}$ of $-20.0 \pm 2\text{‰}$. Ages from *Bison* sp. should be considered minimum ages, given their C_4 -plant-rich diet (Morlan, 1999). Bone collagen from *Homo sapiens* are corrected using an assumed $\delta^{13}\text{C}$ of $-19.0 \pm 2\text{‰}$.

Comparisons between the measured $\delta^{13}\text{C}$ values in the Manitoba dataset show that these values are generally correct (Table 1), and our data fits within that compiled by Morlan (1999). The human collagen in our dataset is slightly more negative than the ‘assumed’, with a mean $\delta^{13}\text{C}$ of -21.4‰ and median of -21.0‰ . The traditional diet in Manitoba was omnivorous, both for terrestrial and aquatic species (Syms, 2018). Though the data are limited, it could mean the four conventional radiocarbon ages on human bone collagen are ~30 to 40 years younger than shown (Gak-5447, S-651, S-743, S-1303). In studies where better precision is needed, it is important to ensure that the ^{13}C ratio is measured and not simply estimated. Syms (2018) suggests that this

ratio is measured for both bone collagen (protein information) and bone apatite (total diet).

Calibrated ages

To compare radiocarbon ages obtained on terrestrial, freshwater and marine organisms within Manitoba, it is necessary to calibrate the ages. It is important to note that radiocarbon ages do not directly equate with calendar years. This is because radiocarbon concentration in the atmosphere varies through time, due to changes in the production rate (de Vries, 1958). As such, calibrations use independently-dated archives such as tree rings, lacustrine and marine sediments, speleothems and corals (Heaton et al., 2020; Reimer et al., 2020). Calibrated ages are accompanied by complex, sometimes multimodal, calibrated age probability distributions that may require stratigraphic information to resolve.

All non-anomalous radiocarbon ages herein (Table 4) were calibrated using the program Calib 8.2 (Stuiver et al., 2020). Ages from terrestrial samples were calibrated using IntCal20 (Reimer et al., 2020) for the northern hemisphere. The marine mollusks were calibrated separately, to remove the reservoir effect from these shells, by using the Marine20 database (Heaton et al., 2020) with an average reservoir off-set (ΔR) of -21 ± 72 for Hudson Bay (Pieńkowski et al., 2022).

The user should note that only the highest-probability age-range is denoted herein ("Cal_BP_2 σ "), with the probability recorded next to the cal. years BP age-range in Table 4 ("Cal_BP_2 σ _probability"). Where the probability is less than 1, the user may want to refer to the probability distributions calculated within Calib 8.2 for alternate age ranges (Stuiver et al., 2020). Calibrated ages are commonly presented in the literature as median ages, with an uncertainty of 1 σ or 2 σ . The 2 σ age-ranges and the median age are included for this purpose. Ages should be presented in publications with all raw data needed for calibration (see Millard, 2014), to be updated by later researchers when new calibration curves are published. Furthermore, the convention is to round the final calculated age to the nearest 10, as the mathematical computations calculate values to a degree that is not precise in reality (Millard, 2014). When using median ages, the user must also include the 2 σ range and the probability (e.g., 8.15 ka cal. years BP, 7960–8340, 100%).

Discussion

Conventional decay-count vs. AMS methods

This GeoFile contains radiocarbon ages determined by conventional decay-count (radiometric) methods and by AMS methods. Both decay-count and AMS methods can provide comparable precision. However, decay-count requires three orders of magnitude more carbon than the AMS method. Hence, historical samples submitted for decay-count methods often used bulk sediment samples, bulk assemblages of macrofossils and/or large pieces of wood. Analysis on bulk samples can lead to

inaccuracies, as these samples may have been contaminated by reworked (older) detritus, overprinted (younger) detritus such as modern rootlets and/or contain different mixtures of materials (e.g., Bayliss and Marshall, 2019). As such, single-specimen radiocarbon samples are considered to provide more accurate results than bulk samples. At sites where organic material has been dated by both conventional and AMS methods, the AMS method is considered more accurate due to single-specimen precision.

Cautions when interpreting ages

The user is reminded that radiocarbon ages are estimates, and should always be considered with regard to other evidence from the site. For both conventional and AMS methods, trace amounts of modern carbon can generate an apparent age that is ultimately incorrect (e.g., Reyes et al., 2020). Reproducibility between (and within) laboratories is also a concern, a problem which seems to increase with older materials (Ward and Clague, 2019). In situations where a hypothesis is based on the result, duplicate radiocarbon ages should be obtained from the same material, and possibly analyzed at different labs (e.g., McMartin et al., 2019). Different pre-treatments may also help to confirm the radiocarbon age (e.g., Bajc et al., 2015). Replicate measurements on different single-specimen samples from the same context or feature can also help (Bayliss and Marshall, 2019). In all cases, other proxies (paleoenvironment, ice-flow dynamics, Heinrich events, other dating methods, cultural context, etc.) should be considered when interpreting a radiocarbon age.

Freshwater reservoir (hard-water) effect

Radiocarbon dating relies on the assumption that organic materials were in equilibrium with the production of ^{14}C in the atmosphere during their lives. Importantly, scientists have learned that organic materials are also affected by inorganic carbon within freshwater environments that overlie both carbonate rocks (Deevey et al., 1954; Andree et al., 1986; MacDonald et al., 1987), and/or lignite, coal and carbonaceous shales (Nambudiri et al., 1980). There may have been different uptake conditions within different lacustrine or fluvial bodies, and within different time periods (Shotton, 1972). Contamination by old-carbon is termed the hard-water effect, and is important in Manitoba because most glacial and postglacial sediments in Manitoba are calcareous (Manitoba Natural Resources and Northern Development, 2022). The hard-water effect is also important when interpreting 'terrestrial' radiocarbon ages obtained from bones of species (birds, humans, canines, bear, wolf, etc.) that may have eaten a marine and/or freshwater diet (Syms, 2018). In Manitoba, the freshwater diet effect ($n = 4$) varies from 220 to 370 ^{14}C years on specimens that are between 440 and 4400 ^{14}C years (Gauthier, 2022). For newer analyses, $^{15}\text{N}/^{14}\text{N}$ ratios should be analyzed to help determine if fish or other aquatic species were part of the diet—if they were, a freshwater (hard-water) correction is needed (Syms, 2018). The column 'Del ^{15}N ' has been added, and it is the intention of the MGS to collect this ratio on all bones and antlers moving forward.

A hard-water correction is typically calculated by comparing the ages of terrestrial and freshwater material collected from the same horizon at the same site. For Manitoba, the hardwater effect ($n = 12$) is temporally variable, with a median of 230 ^{14}C years and values up to at least 880 ^{14}C years (Gauthier, 2022). Eleven of those paired terrestrial-freshwater ages are less than 5000 ^{14}C years old, meaning an accurate hardwater correction for older material is very difficult to determine. To acknowledge the lack of regionally- and temporally-specific data, no hard-water corrections have been made to the data herein. Instead, the notation “FRE?” is added to the conventional and calibrated ages in Table 5 (123 ages), acknowledging the need for future correction as the user finds appropriate. Potential bone samples affected by a freshwater diet (e.g., *Homo sapiens*), and without a paired terrestrial sample, are also included in this table.

Marine reservoir effect

The concentration of carbon isotopes differs between the ocean and the atmosphere, with a ‘reservoir’ in the global oceans that results from trapped old carbon (Arnold, 1957; de Vries, 1958). This reservoir effect results in an apparent radiocarbon age of a marine sample that is different from the true age (when carbon exchange between the organism and the atmosphere would be equal). As such, measured radiocarbon ages from marine samples can’t be directly compared to measured radiocarbon ages from terrestrial samples. Complicating matters, the level of depletion varies due to spatio-temporal differences in the ocean and the atmosphere (Heaton et al., 2020). To account for this, a marine radiocarbon reservoir age is applied to marine organisms used in radiocarbon dating (Stuiver et al., 1986; McNeely et al., 2006; Coulthard et al., 2010). The marine radiocarbon reservoir is calculated using the difference between the global mean ocean reservoir correction and the regional reservoir age, termed the regional reservoir offset ($\Delta R(\theta)$; Heaton et al., 2020). $\Delta R(\theta)$ will remain approximately constant through time, assuming that the regional oceanographic characteristics remain (Stuiver et al., 1986; Stuiver and Braziunas, 1993). Herein, we calibrate the marine radiocarbon ages using the mollusc marine regional reservoir offset $\Delta R(\theta) = -21 \pm 72$ years, for Hudson Bay (Pieńkowski et al., 2022). The regional mollusc marine radiocarbon reservoir (R_r) was calculated using twelve live-collected shells sampled between 1920 and 1954 (pre-bomb).

It should be noted that most Canadian marine radiocarbon reservoir studies involve deposit-feeding molluscs. Recent studies on marine walrus, beluga whale and bowhead whale bones/teeth have determined different reservoir offsets than that of molluscs (Dyke et al., 2019; Pieńkowski et al., 2022), which should caution the user against using mollusc-based corrections for marine mammals. The database herein has one single marine mammal age, which for lack of a better correction, has been tentatively calibrated using the mollusc marine reservoir age (S-521, Rutherford et al., 1973). This calibration is likely wrong.

Marine hard-water effect

Marine materials are affected by inorganic carbon within environments that overlie carbonate-bearing or calcareous rocks or sediment. Open marine water contains dissolved inorganic carbon, which generally masks the hard-water effect. The hard-water effect may be a concern in areas with restricted water circulation, areas of considerable mixing between freshwater and marine waters, overlying highly calcareous substrate, and in areas with high abundance of terrestrial organic matter (Douka et al., 2010). Radiocarbon dating of filter-feeding molluscs will result in avoidance of the hard-water effect relative to deposit-feeding molluscs (England et al., 2013). The database herein contains radiocarbon ages for eight different shell taxa, broken down in Table 3. A ninth taxon, *Portlandia arctica* can be found within the anomalous table, given its proven hard-water effects (England et al., 2013). The twelve *Macoma* sp. radiocarbon ages herein are tagged as ‘Maybe anomalous’ (Table 6), because they are deposit feeders and may need a hard-water correction. Unidentified shells are also classified as ‘maybe anomalous’, given the difficulty of determining feeding behaviour on these ages. Only one site in Manitoba contains radiocarbon ages on bulk samples of both *Hiattella arctica* and *Macoma baltica*. As the *Hiattella arctica* samples (GSC-3367, BGS-791; Nielsen et al., 1986) were collected ~4 m lower than the *Macoma baltica* sample (BGS-797; Morlan et al., 2000), it is impossible to assess a potential hard-water effect—except to note that the upper shells are younger, as expected.

There are very few early Holocene paired terrestrial-marine samples within Hudson Bay, with which to assess a hard-water effect. On Southampton Island, early–mid-Holocene paired terrestrial-marine ages provided a regional marine reservoir age of -630 ± 45 ^{14}C years (Ross et al., 2012) and a reservoir offset (ΔR) age of 263 ± 48 years. That is higher than the ΔR age of 175 ± 89 years for the Foxe basin proposed by Pieńkowski et al. (2022), and suggests the modern correction may need a hard-water

Table 3: Marine shell taxa within the Manitoba radiocarbon database, and their feeding behaviours (after McNeely et al., 2006).

Species	Number of samples	Habitat ¹	Feeding ²
<i>Astarte borealis</i>	1	Infaunal	Suspension
<i>Chlamys islandicus</i>	3	Epifaunal	Suspension
<i>Clinocardium ciliatum</i>	1	Infaunal	Suspension
<i>Hiattella arctica</i>	40	Epifaunal	Suspension
<i>Macoma baltica</i>	2	Infaunal	Deposit
<i>Macoma calcarea</i>	10	Infaunal	Deposit
<i>Mya truncata</i>	4	Epifaunal	Suspension
<i>Mytilus edulis</i>	9	Epifaunal	Suspension
Unidentified	21	?	?

¹ Epifauna are organisms that live upon the surface of sediments. Infauna are organisms that live within sediments.
² Suspension feeders (filter feeders) are animals that feed by straining suspended matter from the water. Deposit feeders are animals that feed by obtaining food particles in the sediment.

correction. Contrastingly, on Baffin Island in the eastern Foxe basin, Vickers et al. (2010) calculated a mean ΔR age of 615 ± 20 years for early–mid Holocene terrestrial-marine pairs. Ross et al. (2012) suggested the difference may be due to underlying geology (the hard-water effect), as northern Southampton Island is granitoid while the Baffin Island sites are calcareous. As such, more local data is needed to ascertain the validity of the corrections applied herein over both time and space.

Infinite and near-infinite ages

Very old samples have such low concentrations of ^{14}C that they cannot be distinguished reliably from the background radiation. Different laboratories set different ‘ages’ as the upper limit, which can vary based on the weight of the material submitted (Pigati et al., 2007); these ages are usually reported as $> xx$ years BP. Regardless of the reported number, the age of these samples was not determined using radiocarbon methods and is interpreted as non-finite (greater than).

As laboratory methods improve over time, the upper boundary of infinite ages has increased. The user is reminded that small amounts of contamination by younger carbon will have large effects near the upper bounds of radiocarbon limits. Contamination can be introduced during burial (diagenetic or modern root-lets), surface weathering, sampling and/or laboratory processing (Pigati et al., 2007). Hence, a lab-accurate age of 48 ka ^{14}C BP may not be a ‘true’ accurate age for that organism. Contamination is more likely for peat (modern roots) or carbonate shells (recrystallization; Douka et al., 2010), than for wood. Recommendations on the ‘validity’ of near-infinite ages range from ~ 40 ka (Walker, 2005) to ~ 35 – 40 ka (Miller and Andrews, 2019)—though the latter paper lacks a source for such a recommendation. In every case, all available geological and site-specific evidence should be considered when interpreting age results, and other proxies (paleoenvironment, ice-flow dynamics, Heinrich events, other dating methods, etc.) should be considered when interpreting the radiocarbon age.

Anomalous and maybe-anomalous ages

A number of radiocarbon ages have been identified as anomalous, either by the author during the course of interpretation, or if the material was bulk organic sediment, lake sediment, or organic mud. The latter are tagged as anomalous given the problems with conventional dating of large mixed samples that may contain older transported detritus and/or younger intruding detritus (e.g., Clayton and Moran, 1982; Grimm et al., 2009; Bayliss and Marshall, 2019). Bulk lake sediment (gyttja) samples also contain nonorganic detritus that contributes to a hard-water effect (see below), which is difficult to correct for given the potential for contamination and re-working. A recent compilation paper by Young et al. (2021) has chosen to reject all bulk lake/soil samples during their analysis—and we advise the same. These rejected samples are retained in the database, and clearly marked ‘yes’ under the column ‘ANOMALOUS’ (263 samples, Table 7). The notation “A” is added to the conventional age, to

further remind the user that the age is likely not valid. The specific reason why each sample was determined to be anomalous is written under column ‘ANOMALOUS_WHY’.

Additional samples are tagged as ‘maybe’ anomalous. This includes stratigraphically inversed (e.g., older above younger) ages and ages near the upper boundaries for radiocarbon dating (>40 ka). Near-finite ages should be confirmed through stratigraphy (including till composition) and re-dating, since they are near the boundary between interstadial marine isotope stage (MIS) 3 (29–57 cal. ka BP; Lisiecki and Raymo, 2005), and MIS 5 (infinite). The notation “M” is added to the conventional age, to further remind the user that the age may not be valid (123 samples, Table 6). The specific reason why each sample was determined to be maybe anomalous is written under column ‘ANOMALOUS_WHY’.

Macrofossil reports

A number of samples were submitted to A. Telka for macrofossil identification. Most of these samples are associated with a macrofossil report, which includes information on different types of plant and animal macrofossils contained within the submitted sediment. These reports are identified with a number under the column “REPORT” and the actual report information is embedded within the “COMMENTS” column of Tables 4–7.

References

- Andree, M., Oeschger, H., Siegenthaler, U., Riesen, T., Moell, M., Ammann, B. and Tobolski, K. 1986: ^{14}C dating of plant macrofossils in lake sediment; *Radiocarbon*, v. 28, no. 2A, p. 411–416.
- Arnold, J.R. 1957: The distribution of carbon-14 in nature; *Tellus*, v. 9, no. 1, p. 28–32.
- Bajc, A.F., Karrow, P.F., Yansa, C.H., Curry, B.B., Nekola, J.C., Seymour, K.L. and Mackie, G.L. 2015: Geology and paleoecology of a Middle Wisconsin fossil occurrence in Zorra Township, southwestern Ontario, Canada; *Canadian Journal of Earth Sciences*, v. 52, no. 6, p. 386–404, URL <<https://doi.org/10.1139/cjes-2015-0005>>.
- Bayliss, A. and Marshall, P. 2019: Confessions of a serial polygamist: the reality of radiocarbon reproducibility in archaeological samples; *Radiocarbon*, v. 61, no. 5, p. 1143–1158, URL <<https://doi.org/10.1017/RDC.2019.55>>.
- Clayton, L. and Moran, S.R. 1982: Chronology of late Wisconsinan glaciations in middle North America; *Quaternary Science Reviews*, v. 1, p. 55–82.
- Coulthard, R.D., Furze, M.F.A., Pieńkowski, A.J., Nixon, F.C. and England, J.H. 2010: New marine ΔR values for Arctic Canada; *Quaternary Geochronology*, v. 5, no. 4, p. 419–434.
- Dalton, A.S., Margold, M., Stokes, C.R., Tarasov, L., Dyke, A.S., Adams, R.S., Allard, S., Arends, H., Atkinson, N., Attig, J.W., Barnett, P.J., Barnett, R.L., Batterson, M., Bernatchez, P., Borns, H.W., Breckenridge, A., Briner, J.P., Brouard, E., Campbell, J.E., Carlson, A.E. et al. 2020: An updated radiocarbon-based ice margin chronology for the last deglaciation of the North American Ice Sheet Complex; *Quaternary Science Reviews*, v. 234, art. 106223, URL <<https://doi.org/10.1016/j.quascirev.2020.106223>>.
- de Vries, H. 1958: Variations in concentration of radiocarbon with time and location on earth; *Koninklijke Nederlandse Akademie van Wetenschappen, Proc. Ser. B*, v. 61, no. 2, p. 94–102.

- Deevey, E.S., Jr., Gross, M.S., Hutchinson, G.E. and Kraybill, H.L. 1954: The natural C^{14} contents of materials from hard-water lakes; *PNAS*, v. 40, no. 5, p. 285–288, URL <<https://doi.org/10.1073/pnas.40.5.285>>.
- Douka, K., Higham, T.F.G. and Hedges, R.E.M. 2010: Radiocarbon dating of shell carbonates: old problems and new solutions; *Munibe Suplemento*, v. 31, p. 18–27.
- Dyke, A.S., Saville, J.M., Szpak, P., Southon, J.R., Howse, L., Desrosiers, P.M. and Kotar, K. 2019: An assessment of marine reservoir corrections for radiocarbon dates on walrus from the Foxe Basin region of Arctic Canada; *Radiocarbon*, v. 61, no. 1, p. 67–81, URL <<https://doi.org/10.1017/RDC.2018.50>>.
- England, J., Dyke, A.S., Coulthard, R.D., McNeely, R. and Aitken, A. 2012: The exaggerated radiocarbon age of deposit-feeding molluscs in calcareous environments; *Boreas*, v. 42, no. 2, p. 362–373, URL <<https://doi.org/10.1111/j.1502-3885.2012.00256.x>>.
- Fritz, P. and Poplawski, S. 1974: ^{18}O and ^{13}C in the shells of freshwater molluscs and their environments; *Earth and Planetary Science Letters*, v. 24, no. 1, p. 91–98.
- Gauthier, M.S. 2021: Manitoba radiocarbon ages: update; Manitoba Agriculture and Resource Development, Manitoba Geological Survey, Open File OF2021-1, 7 p. plus 2 appendices.
- Gauthier, M.S. 2022: Using radiocarbon ages on organics affected by freshwater—a geologic and archaeologic update on the freshwater reservoir ages and freshwater diet effect in Manitoba, Canada; *Radiocarbon*, v. 64, no. 2, p. 253–264, URL <<https://doi.org/10.1017/RDC.2022.30>>.
- Gauthier, M.S., Santucci, A. and Keller, G.R. 2022: Digital compilation of surficial point and line features for Manitoba, including ice-flow data; Manitoba Natural Resources and Northern Development, Manitoba Geological Survey, GeoFile 1-2022.
- Government of Manitoba 2020: Southern Manitoba LIDAR data; URL <https://mli2.gov.mb.ca/dems/index_external_lidar.html> [January 2020].
- Grimm, E.C., Maher, L.J., Jr. and Nelson, D.M. 2009: The magnitude of error in conventional bulk-sediment radiocarbon dates from central North America; *Quaternary Research*, v. 72, no. 2, p. 301–308.
- Heaton, T.J., Kohler, P., Butzin, M., Bard, E., Reimer, R.W., Austin, W. E.N., Bronk Ramsey, C., Grootes, P.M., Hughen, K.A., Kromer, B., Reimer, P.J., Adkins, J., Burke, A., Cook, M.S., Olsen, J. and Skinner, L.C. 2020: Marine20—the marine radiocarbon age calibration curve (0–55,000 cal BP); *Radiocarbon*, v. 62, no. 4, p. 779–820, URL <<https://doi.org/10.1017/RDC.2020.68>>.
- Lisiecki, L.E. and Raymo, M.E. 2005: A Pliocene-Pleistocene stack of 57 globally distributed benthic $\delta^{18}O$ records; *Paleoceanography and Paleoclimatology*, v. 20, no. 1, art. PA1003.
- MacDonald, G.M., Beukens, R.P., Kieser, W.E. and Vitt, D.H. 1987: Comparative radiocarbon dating of terrestrial plant macrofossils and aquatic moss from the “ice-free corridor” of western Canada; *Geology*, v. 15, no. 9, p. 837–840, URL <[https://doi.org/10.1130/0091-7613\(1987\)15<837:CRDOTP>2.0.CO;2](https://doi.org/10.1130/0091-7613(1987)15<837:CRDOTP>2.0.CO;2)>.
- Manitoba Natural Resources and Northern Development 2022: Manitoba carbonate dispersal analyses in till; Manitoba Natural Resources and Northern Development, Manitoba Geological Survey, URL <<http://manitoba.ca/iem/geo/surficial/carbonate.html>> [July 2022].
- Martindale, A., Morlan, R., Betts, M., Blake, M., Gajewski, K., Chaput, M., Mason, A. and Vermeersch, P. 2016: Canadian archaeological radiocarbon database (CARD 2.1); URL <<https://www.canadianarchaeology.ca>> [December 2020].
- Marty, J. and Myrbo, A. 2014: Radiocarbon dating suitability of aquatic plant macrofossils; *Journal of Paleolimnology*, v. 52, p. 435–443.
- McMartin, I., Campbell, J.E. and Dredge, L.A. 2019: Middle Wisconsinan marine shells near Repulse Bay, Nunavut, Canada: implications for marine isotope stage 3 ice-free conditions and Laurentide Ice Sheet dynamics in north-west Hudson Bay; *Journal of Quaternary Science*, v. 34, no. 1, p. 64–75, URL <<https://doi.org/10.1002/jqs.3081>>.
- McNeely, R. and Brennan, J. 2005: Geological Survey of Canada revised shell dates; Geological Survey of Canada, Open File 5019, 530 p.
- McNeely, R., Dyke, A.S. and Southon, J.R. 2006: Canadian marine reservoir ages, preliminary data assessment; Geological Survey of Canada, Open File 5049, 3 p.
- Millard, A.R. 2014: Conventions for reporting radiocarbon determinations; *Radiocarbon*, v. 56, no. 2, p. 555–559, URL <<https://doi.org/10.2458/56.17455>>.
- Miller, G.H. and Andrews, J.T. 2019: Hudson Bay was not deglaciated during MIS-3; *Quaternary Science Reviews*, v. 225, no. 1, art. 105944, URL <<https://doi.org/10.1016/j.quascirev.2019.105944>>.
- Morlan, R. 1999: Canadian archaeological radiocarbon database: establishing conventional ages; *Canadian Journal of Archaeology*, v. 23, no. 1/2, p. 3–10, URL <<https://www.jstor.org/stable/41103370>> [June 2023].
- Morlan, R.E., McNeely, R. and Nielsen, E. 2000: Manitoba radiocarbon dates; Manitoba Industry, Trade and Mines, Manitoba Geological Survey, Open File Report OF2000-1, 198 p.
- Nambudiri, E.M.V., Teller, J.T. and Last, W.M. 1980: Pre-Quaternary microfossils—a guide to errors in radiocarbon dating; *Geology*, v. 8, no. 3, p. 123–126, URL <[https://doi.org/10.1130/0091-7613\(1980\)8<123:PMGTET>2.0.CO;2](https://doi.org/10.1130/0091-7613(1980)8<123:PMGTET>2.0.CO;2)>.
- Nielsen, E., Morgan, A.V., Morgan, A., Mott, R.J., Rutter, N.W. and Causse, C. 1986: Stratigraphy, paleoecology and glacial history of the Gillam area, Manitoba; *Canadian Journal of Earth Sciences*, v. 23, no. 11, p. 1641–1661.
- Philippson, B. 2013: The freshwater reservoir effect in radiocarbon dating; *Heritage Science*, v. 1, no. 24, 19 p., URL <<https://heritagesciencejournal.springeropen.com/articles/10.1186/2050-7445-1-24>> [June 2023].
- Pieńkowski, A.J., Coulthard, R.D. and Furze, M.F. A. 2022: Revised marine reservoir offset (ΔR) values for molluscs and marine mammals from Arctic North America; *Boreas*, v. 52, no. 2, p. 145–167, URL <<https://doi.org/10.1111/bor.12606>>.
- Pigati, J.S., Quade, J., Wilson, J., Jull, A.J.T. and Lifton, N.A. 2007: Development of low-background vacuum extraction and graphitization systems for ^{14}C dating of old (40–60ka) samples; *Quaternary International*, v. 166, no. 1, p. 4–14.
- Reimer, P.J., Austin, W.E.N., Bard, E., Bayliss, A., Blackwell, P.G., Bronk Ramsey, C., Butzin, M., Cheng, H., Edwards, R.L., Friedrich, M., Grootes, P.M., Guilderson, T.P., Hajdas, I., Heaton, T.J., Hogg, A.G., Hughen, K.A., Kromer, B., Manning, S., Muscheler, R., Palmer, J.G. et al. 2020: The IntCal20 northern hemisphere radiocarbon age calibration curve (0–55 cal kBP); *Radiocarbon*, v. 62, no. 4, p. 725–757, URL <<https://doi.org/10.1017/RDC.2020.41>>.
- Reyes, A.V., Dillman, T., Kennedy, K., Froese, D., Beaudoin, A.B. and Paulen, R.C. 2020: Legacy radiocarbon ages and the MIS 3 dating game: a cautionary tale from re-dating of pre-LGM sites in western Canada; Geological Society of America, GSA 2020 Connects Online virtual event, October 26–30, 2020, Abstracts with Programs, v. 52, no. 6, p. 1, paper no. 56-3, URL <<https://doi.org/10.1130/abs/2020AM-360064>>.
- Ross, M., Utting, D.J., Lajeunesse, P. and Kosar, K.G.A. 2012: Early Holocene deglaciation of northern Hudson Bay and Foxe Channel constrained by new radiocarbon ages and marine reservoir correction; *Quaternary Research*, v. 78, no. 1, p. 82–94.

- Rutherford, A.A., Wittenberg, J. and McCallum, K.J. 1973: University of Saskatchewan radiocarbon dates VI; *Radiocarbon*, v. 15, no. 1, p. 193–211.
- Shotton, F.W. 1972: An example of hard-water error in radiocarbon dating of vegetable matter; *Nature*, v. 240, p. 460–461, URL <<https://doi.org/10.1038/240460a0>>.
- Stuiver, M. and Braziunas, T.F. 1993: Modeling atmospheric ^{14}C influences and ^{14}C ages of marine samples to 10,000 BC; *Radiocarbon*, v. 35, no. 1, p. 137–189, URL <<https://doi.org/10.1017/S0033822200013874>>.
- Stuiver, M. and Polach, H.A. 1977: Discussion reporting of ^{14}C data; *Radiocarbon*, v. 19, no. 3, p. 355–363.
- Stuiver, M., Pearson, G.W. and Braziunas, T. 1986: Radiocarbon age calibration of marine samples back to 9000 cal yr BP; *Radiocarbon*, v. 28, no. 2B, p. 980–1021.
- Stuiver, M., Reimer, P.J. and Reimer, R. 2020: CALIB 8.2 [WWW program]; URL <<http://calib.org>> [December 2020].
- Syms, E.L. 2018: Discovering the fresh water reservoir effect in the northern boreal forests: refining the temporal framework; *Ontario Archaeology*, no. 98, p. 48–65.
- Teller, J.T. 1980: Radiocarbon dates in Manitoba; Manitoba Department of Energy and Mines, Mineral Resources Division, Geological Report GR80-4, 61 p.
- Trumbore, S.E. 2000: Radiocarbon geochronology; in *Quaternary Geochronology: methods and applications*, J.S. Noller, J.M. Sowers and W.R. Lettis (ed.), American Geophysical Union, Washington, D.C., p. 41–60.
- U.S. Geological Survey 2014: USGS EROS archive - digital elevation - Shuttle Radar Topography Mission (SRTM) 1 arc-second global; U.S. Geological Survey, 30 m cell, zipped hgt format, URL <<https://doi.org/10.5066/F7PR7TFT>>.
- Vickers, K.J., Ward, B.C., Utting, D.J. and Telka, A.M. 2010: Deglacial reservoir age and implications, Foxe Peninsula, Baffin Island; *Journal of Quaternary Science*, v. 25, p. 1388–1346, URL <<https://doi.org/10.1002/jqs.1419>>.
- Walker, M. 2005: *Quaternary dating methods*; John Wiley & Sons, Ltd., Chichester, West Sussex, United Kingdom, 304 p.
- Ward, B.C. and Clague, J. 2019: A blind comparison of radiocarbon labs; European Geosciences Union, EGU General Assembly 2019, April 7–12, 2019, Vienna, Austria, Geophysical Research Abstracts, v. 21, EGU2019-12003.
- Young, J.M., Reyes, A.V. and Froese, D.G. 2021: Assessing the ages of the Moorhead and Emerson phases of glacial Lake Agassiz and their temporal connection to the Younger Dryas cold reversal; *Quaternary Science Reviews*, v. 251, art. 106714, URL <<https://doi.org/10.1016/j.quascirev.2020.106714>>.